

Provided for non-commercial research and educational use.
Not for reproduction, distribution or commercial use.

Serdica

Mathematical Journal

Сердика

Математическо списание

The attached copy is furnished for non-commercial research and education use only.
Authors are permitted to post this version of the article to their personal websites or institutional repositories and to share with other researchers in the form of electronic reprints.
Other uses, including reproduction and distribution, or selling or licensing copies, or posting to third party websites are prohibited.

For further information on
Serdica Mathematical Journal
which is the new series of
Serdica Bulgaricae Mathematicae Publicationes
visit the website of the journal <http://www.math.bas.bg/~serdica>
or contact: Editorial Office
Serdica Mathematical Journal
Institute of Mathematics and Informatics
Bulgarian Academy of Sciences
Telephone: (+359-2)9792818, FAX:(+359-2)971-36-49
e-mail: serdica@math.bas.bg

OSCILLATION CRITERIA FOR NONLINEAR DIFFERENTIAL EQUATIONS OF SECOND ORDER WITH DAMPING TERM

E. M. Elabbasy, W. W. Elhaddad

Communicated by E. I. Horozov

ABSTRACT. Some new criteria for the oscillation of all solutions of second order differential equations of the form

$$\frac{d}{dt} \left(r(t)\psi(x) \left| \frac{dx}{dt} \right|^{\alpha-2} \frac{dx}{dt} \right) + p(t)\varphi \left(|x|^{\alpha-2} x, r(t)\psi(x) \left| \frac{dx}{dt} \right|^{\alpha-2} \frac{dx}{dt} \right) + q(t) |x|^{\alpha-2} x = 0,$$

and the more general equation

$$\frac{d}{dt} \left(r(t)\psi(x) \left| \frac{dx}{dt} \right|^{\alpha-2} \frac{dx}{dt} \right) + p(t)\varphi \left(g(x), r(t)\psi(x) \left| \frac{dx}{dt} \right|^{\alpha-2} \frac{dx}{dt} \right) + q(t)g(x) = 0,$$

are established. our results generalize and extend some known oscillation criterain in the literature.

2000 *Mathematics Subject Classification*: 34C10, 34C15.

Key words: Oscillation, second order nonlinear differential equation.

1. Introduction. We are concerned with the oscillation of solutions of second order differential equations with damping of the following form

$$(E_1) \quad \frac{d}{dt} \left(r(t)\psi(x) \left| \frac{dx}{dt} \right|^{\alpha-2} \frac{dx}{dt} \right) + p(t)\varphi \left(|x|^{\alpha-2} x, r(t)\psi(x) \left| \frac{dx}{dt} \right|^{\alpha-2} \frac{dx}{dt} \right) \\ + q(t) |x|^{\alpha-2} x = 0,$$

and the more general equation

$$(E_2) \quad \frac{d}{dt} \left(r(t)\psi(x) \left| \frac{dx}{dt} \right|^{\alpha-2} \frac{dx}{dt} \right) + p(t)\varphi \left(g(x), r(t)\psi(x) \left| \frac{dx}{dt} \right|^{\alpha-2} \frac{dx}{dt} \right) \\ + q(t)g(x) = 0,$$

where $r \in C[[t_0, \infty), \mathbf{R}^+]$, $p \in C[[t_0, \infty), [0, \infty)]$, $q \in C[[t_0, \infty), \mathbf{R}]$, $\psi \in C[\mathbf{R}, \mathbf{R}^+]$ and $g \in C^1[\mathbf{R}, \mathbf{R}]$ such that $xg(x) > 0$ for $x \neq 0$ and $\frac{d}{dx}g(x) > 0$ for $x \neq 0$. φ is defined and continuous on $\mathbf{R} \times \mathbf{R} - \{0\}$ with $u\varphi(u, v) > 0$ for $uv \neq 0$ and $\varphi(\lambda u, \lambda v) = \lambda\varphi(u, v)$ for $0 < \lambda < \infty$ and $(u, v) \in \mathbf{R} \times \mathbf{R} - \{0\}$.

By the oscillation of equation (E_1) [(E_2)], we mean a function $x \in C^1([T_x, \infty), \mathbf{R})$ for some $T_x \geq t_0$, which has the property that $r(t)\psi(x) \left| \frac{dx}{dt} \right|^{\alpha-2} \frac{dx}{dt} \in C^1([T_x, \infty), \mathbf{R})$ and satisfies equation (E_1) [(E_2)] on $[T_x, \infty)$.

A solution of equation (E_1) [(E_2)] is called oscillatory if it has arbitrarily large zeros otherwise, it is called nonoscillatory. Finally, equation (E_1) [(E_2)] is called oscillatory if all its solutions are oscillatory.

In Section 2 we provide sufficient conditions for the oscillation of all solutions of (E_1) . Several particular cases of (E_1) have been discussed in the literature. To cite a few examples, the differential equation

$$(E_3) \quad \frac{d}{dt} \left(r(t) \left| \frac{dx}{dt} \right|^{\alpha-2} \frac{dx}{dt} \right) + q(t) |x|^{\alpha-1} x = 0$$

has been studied by Hsu and yeh [2] Kusano and Naito [4] Kusano, Yoshida [5], Li and Yeh [6], [7], [8], [9], [10] and Lian, Yeh and Li [11]. A more general equation than (E_3)

$$(E_4) \quad \frac{d}{dt} \left(r(t)\psi(x) \left| \frac{dx}{dt} \right|^{\alpha-2} \frac{dx}{dt} \right) + q(t) |x|^{\alpha-1} x = 0$$

has been considered by Ayanlar and Tiryaki [1] and Wu et al. [16]. Our results include, as special cases, known oscillation theorems for (E_3) and (E_4) . In particular, we extend and improve the results obtained in [12], [16] and [14].

In Section 3 we will establish some oscillation criteria for equation (E_2) . Several particular cases of (E_2) have been discussed in the literature. The differential equation

$$(E_5) \quad \frac{d}{dt} \left(r(t)\psi(x) \left| \frac{dx}{dt} \right|^{\alpha-2} \frac{dx}{dt} \right) + q(t)g(x) = 0,$$

established by Manojlovic [13]. Wong and Agarwal [15] considered a special case of this equation as

$$(E_6) \quad \frac{d}{dt} \left(r(t) \left| \frac{dx}{dt} \right|^{\alpha-2} \frac{dx}{dt} \right) + q(t)g(x) = 0.$$

Our results in this section generalize and improve Manojlovic [13].

2. Oscillation results for (E_1) . In order to discuss our main results, we need the following well-known inequality which is due to Hardy et al. [3, Theorem 41].

Lemma 1. *If X and Y are nonnegative, then*

$$X^\lambda + (\lambda - 1)Y^\lambda - \lambda XY^{\lambda-1} \geq 0, \quad \lambda > 1,$$

where equality holds if and only if $X = Y$.

Theorem 1. *Suppose that*

$$(1) \quad \varphi(1, z) \geq z \quad \text{for all } z \neq 0,$$

$$(2) \quad 0 < \psi(x) \leq \gamma \quad \text{for all } x,$$

and there exist differentiable functions

$$k, \rho : [t_0, \infty) \rightarrow (0, \infty),$$

and the continuous function

$$H : D \equiv \{(t, s) : t \geq s \geq t_0\} \rightarrow \mathbf{R} \quad \text{and} \quad h : D_0 \equiv \{(t, s) : t > s \geq t_0\} \rightarrow \mathbf{R},$$

where H has a continuous nonpositive partial derivative on D with respect to the second variable such that

$$H(t, t) = 0 \text{ for } t \geq t_0, \quad H(t, s) > 0 \text{ for } t > s \geq t_0,$$

and

$$h(t, s) = -\frac{\partial}{\partial s} (H(t, s)k(s)) - \left(\frac{\frac{d}{ds}\rho(s)}{\rho(s)} - p(s) \right) H(t, s)k(s) \text{ for all } (t, s) \in D_0.$$

Then equation (E_1) is oscillatory if

$$(3) \quad \limsup_{t \rightarrow \infty} \frac{1}{H(t, t_0)} \int_{t_0}^t \left[H(t, s)\rho(s)k(s)q(s) - \frac{\gamma\rho(s)r(s)|h(t, s)|^\alpha}{\alpha^\alpha(H(t, s)k(s))^{\alpha-1}} \right] ds = \infty.$$

Proof. On the contrary we assume that (E_1) has a nonoscillatory solution $x(t)$. We suppose without loss of generality that $x(t) > 0$ for all $t \in [t_0, \infty)$. We define the function $\omega(t)$ as

$$\omega(t) = \rho(t) \frac{r(t)\psi(x) \left| \frac{dx}{dt} \right|^{\alpha-2} \frac{dx}{dt}}{|x|^{\alpha-2} x} \text{ for } t \geq t_0.$$

Thus,

$$\frac{d}{dt}\omega(t) = \frac{\frac{d}{dt}\rho(t)}{\rho(t)}\omega(t) + \rho(t) \frac{\frac{d}{dt} \left(r(t)\psi(x) \left| \frac{dx}{dt} \right|^{\alpha-2} \frac{dx}{dt} \right)}{|x|^{\alpha-2} x} - (\alpha-1) \frac{\rho(t)r(t)\psi(x) \left| \frac{dx}{dt} \right|^\alpha}{|x|^\alpha}.$$

This and equation (E_1) imply

$$\frac{d}{dt}\omega(t) \leq \frac{\frac{d}{dt}\rho(t)}{\rho(t)}\omega(t) - \rho(t)[q(t) + p(t)\varphi(1, \frac{\omega(t)}{\rho(t)})] - (\alpha-1)[\gamma\rho(t)r(t)]^{\frac{-1}{\alpha-1}} |\omega(t)|^{\frac{\alpha}{\alpha-1}}.$$

From (1) we obtain

$$\frac{d}{dt}\omega(t) \leq \frac{\frac{d}{dt}\rho(t)}{\rho(t)}\omega(t) - \rho(t)q(t) - p(t)\omega(t) - (\alpha-1)[\gamma\rho(t)r(t)]^{\frac{-1}{\alpha-1}} |\omega(t)|^{\frac{\alpha}{\alpha-1}}.$$

Multiply the above inequality by $H(t, s)k(s)$ and integrate from T to t we obtain

$$\int_T^t H(t, s)k(s)\rho(s)q(s)ds \leq \int_T^t H(t, s)k(s) \left(\frac{\frac{d}{ds}\rho(s)}{\rho(s)} - p(s) \right) \omega(s)ds$$

$$- \int_T^t H(t, s)k(s)\frac{d}{ds}\omega(s)ds - (\alpha - 1) \int_T^t H(t, s)k(s)[\gamma\rho(s)r(s)]^{\frac{-1}{\alpha-1}} |\omega(s)|^{\frac{\alpha}{\alpha-1}} ds.$$

Since

$$- \int_T^t H(t, s)k(s)\frac{d}{ds}\omega(s)ds = H(t, T)k(T)\omega(T) + \int_T^t \frac{\partial}{\partial s}(H(t, s)k(s))\omega(s)ds,$$

the previous inequality becomes

$$\int_T^t H(t, s)k(s)\rho(s)q(s)ds \leq H(t, T)k(T)\omega(T) - \int_T^t h(t, s)\omega(s)ds$$

$$- (\alpha - 1) \int_T^t H(t, s)k(s)[\gamma\rho(s)r(s)]^{-1/(\alpha-1)} |\omega(s)|^{\alpha/(\alpha-1)} ds.$$

Hence we have

$$\int_T^t H(t, s)k(s)\rho(s)q(s)ds \leq H(t, T)k(T)\omega(T) + \int_T^t |h(t, s)| |\omega(s)| ds$$

$$(4) \quad - (\alpha - 1) \int_T^t H(t, s)k(s)[\gamma\rho(s)r(s)]^{-1/(\alpha-1)} |\omega(s)|^{\alpha/(\alpha-1)} ds.$$

Define

$$X = [\gamma\rho(s)r(s)]^{1/\alpha} \frac{h(t, s)}{\alpha (H(t, s)k(s))^{\alpha/(\alpha-1)}},$$

$$Y = \left(H(t, s)k(s)[\gamma\rho(s)r(s)]^{-1/(\alpha-1)} |\omega(s)|^{\alpha/(\alpha-1)} \right)^{1/\alpha}.$$

Since $\alpha > 1$, by Lemma 1 we obtain

$$|h(t, s)| |\omega(s)| - \frac{(\alpha - 1)H(t, s)k(s)}{[\gamma\rho(s)r(s)]^{1/(\alpha-1)}} |\omega(s)|^{\alpha/(\alpha-1)} \leq \frac{\gamma\rho(s)r(s) |h(t, s)|^\alpha}{\alpha^\alpha (H(t, s)k(s))^{\alpha-1}},$$

for all $t > s \geq T$. Moreover, by (4), we also have for every $t \geq T$,

$$\int_T^t H(t, s)k(s)\rho(s)q(s)ds \leq H(t, T)k(T)\omega(T) + \int_T^t \frac{\gamma\rho(s)r(s)|h(t, s)|^\alpha}{\alpha^\alpha (H(t, s)k(s))^{\alpha-1}} ds,$$

or

$$(5) \int_T^t \left[H(t, s)k(s)\rho(s)q(s) - \frac{\gamma\rho(s)r(s)|h(t, s)|^\alpha}{\alpha^\alpha (H(t, s)k(s))^{\alpha-1}} \right] ds \leq H(t, T)k(T)\omega(T) \\ \leq H(t, T)k(T) |\omega(T)| \\ \leq H(t, t_0)k(T) |\omega(T)|.$$

We use the above inequality for $T = T_0$ to obtain

$$\int_{T_0}^t \left[H(t, s)k(s)\rho(s)q(s) - \frac{\gamma\rho(s)r(s)|h(t, s)|^\alpha}{\alpha^\alpha (H(t, s)k(s))^{\alpha-1}} \right] ds \leq H(t, t_0)k(T_0) |\omega(T_0)|.$$

Therefore,

$$\int_{t_0}^t \left[H(t, s)k(s)\rho(s)q(s) - \frac{\gamma\rho(s)r(s)|h(t, s)|^\alpha}{\alpha^\alpha (H(t, s)k(s))^{\alpha-1}} \right] ds \\ = \int_{t_0}^{T_0} \left[H(t, s)k(s)\rho(s)q(s) - \frac{\gamma\rho(s)r(s)|h(t, s)|^\alpha}{\alpha^\alpha (H(t, s)k(s))^{\alpha-1}} \right] ds \\ + \int_{T_0}^t \left[H(t, s)k(s)\rho(s)q(s) - \frac{\gamma\rho(s)r(s)|h(t, s)|^\alpha}{\alpha^\alpha} \right] ds.$$

Hence for every $t \geq t_0$ we have

$$\int_{t_0}^t \left[H(t, s)k(s)\rho(s)q(s) - \frac{\gamma\rho(s)r(s)|h(t, s)|^\alpha}{\alpha^\alpha (H(t, s)k(s))^{\alpha-1}} \right] ds \\ \leq H(t, t_0) \int_{t_0}^{T_0} k(s)\rho(s) |q(s)| ds + H(t, t_0)k(T_0) |\omega(T_0)| \\ = H(t, t_0) \left\{ \int_{t_0}^{T_0} k(s)\rho(s) |q(s)| ds + k(T_0) |\omega(T_0)| \right\}.$$

This gives

$$\limsup_{t \rightarrow \infty} \frac{1}{H(t, t_0)} \int_{t_0}^t \left[H(t, s)\rho(s)k(s)q(s) - \frac{\gamma\rho(s)r(s)|h(t, s)|^\alpha}{\alpha^\alpha (H(t, s)k(s))^{\alpha-1}} \right] ds$$

$$\leq \left\{ \int_{t_0}^{T_0} k(s)\rho(s) |q(s)| ds + k(T_0) |\omega(T_0)| \right\} < \infty$$

which contradicts the assumption (3). This completes the proof. \square

Corollary 1. *If the condition (3) is replaced by the condition*

$$\limsup_{t \rightarrow \infty} \frac{1}{H(t, t_0)} \int_{t_0}^t H(t, s)\rho(s)k(s)q(s)ds = \infty,$$

and

$$\limsup_{t \rightarrow \infty} \frac{1}{H(t, t_0)} \int_{t_0}^t \frac{\rho(s)r(s) |h(t, s)|^\alpha}{(H(t, s)k(s))^{\alpha-1}} ds < \infty,$$

then the conclusion of Theorem 1 remains valid.

Theorem 2. *Suppose that (1) and (2) hold, and let the functions H, h, ρ and k be the same as in Theorem 1. Moreover, assume that*

$$(6) \quad 0 < \inf_{s \geq t_0} \left[\liminf_{t \rightarrow \infty} \frac{H(t, s)}{H(t, t_0)} \right] \leq \infty,$$

and

$$(7) \quad \limsup_{t \rightarrow \infty} \frac{1}{H(t, t_0)} \int_{t_0}^t \frac{\rho(s)r(s) |h(t, s)|^\alpha}{(H(t, s)k(s))^{\alpha-1}} < \infty,$$

hold. If there exists a function $\Omega \in C([t_0, \infty), \mathbf{R})$ such that

$$(8) \quad \limsup_{t \rightarrow \infty} \int_{t_0}^t \frac{\Omega_+^{\alpha/(\alpha-1)}(s)}{(k(s)\rho(s)r(s))^{1/(\alpha-1)}} = \infty,$$

and for every $T \geq t_0$,

$$(9) \quad \limsup_{t \rightarrow \infty} \frac{1}{H(t, T)} \int_T^t \left[H(t, s)\rho(s)k(s)q(s) - \frac{\gamma\rho(s)r(s) |h(t, s)|^\alpha}{\alpha^\alpha (H(t, s)k(s))^{\alpha-1}} \right] ds \geq \Omega(T),$$

where $\Omega_+(t) = \max\{\Omega(t), 0\}$ for $t \geq t_0$, then equation (E_1) is oscillatory.

Proof. On the contrary we assume that (E_1) has a nonoscillatory solution $x(t)$. We suppose without loss of generality that $x(t) > 0$ for all $t \in [t_0, \infty)$. Defining $\omega(t)$ as in the proof of Theorem 1, we obtain (4) and (5). Then, for $t > T \geq t_0$ we have

$$\limsup_{t \rightarrow \infty} \frac{1}{H(t, T)} \int_T^t \left[H(t, s)k(s)\rho(s)q(s) - \frac{\gamma\rho(s)r(s) |h(t, s)|^\alpha}{\alpha^\alpha (H(t, s)k(s))^{\alpha-1}} \right] ds \leq k(T)\omega(T).$$

Thus, by (9) we have

$$(10) \quad \Omega(T) \leq k(T)\omega(T) \quad \text{for all } T \geq T_0,$$

and

$$(11) \quad \limsup_{t \rightarrow \infty} \frac{1}{H(t, T_0)} \int_{T_0}^t H(t, s)\rho(s)k(s)q(s)ds \geq \Omega(T_0).$$

Let

$$F(t) = \frac{1}{H(t, T_0)} \int_{T_0}^t |h(t, s)| |\omega(s)| ds,$$

$$G(t) = \frac{(\alpha - 1)}{H(t, T_0)} \int_{T_0}^t H(t, s)k(s)[\rho(s)r(s)]^{\frac{-1}{\alpha-1}} |\omega(s)|^{\frac{\alpha}{\alpha-1}} ds,$$

for $t > T_0$. Then by (4) and (11), we get that

$$(12) \quad \liminf_{t \rightarrow \infty} |G(t) - F(t)| \leq k(T_0)\omega(T_0) - \limsup_{t \rightarrow \infty} \frac{1}{H(t, T_0)} \int_{T_0}^t H(t, s)\rho(s)k(s)q(s)ds \leq k(T_0)\omega(T_0) - \Omega(T_0) < \infty.$$

Now, we claim that

$$(13) \quad \int_{T_0}^{\infty} k(s) \frac{|\omega(s)|^{\alpha/(\alpha-1)}}{[\rho(s)r(s)]^{1/(\alpha-1)}} < \infty.$$

Suppose to the contrary that

$$(14) \quad \int_{T_0}^{\infty} k(s) \frac{|\omega(s)|^{\alpha/(\alpha-1)}}{[\rho(s)r(s)]^{1/(\alpha-1)}} = \infty.$$

By (6), there is a positive constant η satisfying

$$(15) \quad \inf_{s \geq t_0} \left[\liminf_{t \rightarrow \infty} \frac{H(t, s)}{H(t, t_0)} \right] > \eta.$$

On the other hand, by (14) for any positive number μ there exists a $T_1 > T_0$ such that

$$\int_{T_0}^t k(s) \frac{|\omega(s)|^{\alpha/(\alpha-1)}}{[\rho(s)r(s)]^{1/(\alpha-1)}} ds \geq \frac{\mu}{(\alpha - 1)\eta} \quad \text{for all } t \geq T_1.$$

So for all $t \geq T_1$

$$\begin{aligned}
 G(t) &= \frac{(\alpha - 1)}{H(t, T_0)} \int_{T_0}^t H(t, s) d \left[\int_{T_0}^s k(u) \frac{|\omega(u)|^{\alpha/(\alpha-1)}}{[\rho(u)r(u)]^{1/(\alpha-1)}} du \right] \\
 &= \frac{(\alpha - 1)}{H(t, T_0)} \int_{T_0}^t \left[\frac{-\partial H(t, s)}{\partial s} \right] d \left[\int_{T_0}^s k(u) \frac{|\omega(u)|^{\alpha/(\alpha-1)}}{[\rho(u)r(u)]^{1/(\alpha-1)}} du \right] ds \\
 &\geq \frac{(\alpha - 1)}{H(t, T_0)} \int_{T_1}^t \left[\frac{-\partial H(t, s)}{\partial s} \right] d \left[\int_{T_0}^s k(u) \frac{|\omega(u)|^{\alpha/(\alpha-1)}}{[\rho(u)r(u)]^{1/(\alpha-1)}} du \right] ds \\
 (16) \quad &\geq \frac{\mu}{(\alpha - 1)\eta} \frac{(\alpha - 1)}{H(t, T_0)} \int_{T_1}^t \left[\frac{-\partial H(t, s)}{\partial s} \right] ds = \frac{\mu H(t, T_1)}{\eta H(t, T_0)}.
 \end{aligned}$$

From (15) we have

$$\liminf_{t \rightarrow \infty} \frac{H(t, T_1)}{H(t, t_0)} > \eta > 0.$$

So there exists $T_2 \geq T_1$ such that $\frac{H(t, T_1)}{H(t, t_0)} \geq \eta$ for all $t \geq T_2$. Therefore by (16) $G(t) \geq \mu$ for all $t \geq T_2$, and since μ is arbitrary constant, we conclude that

$$(17) \quad \lim_{t \rightarrow \infty} G(t) = \infty.$$

Next, consider a sequence $\{t_n\}_{n=1}^\infty$ in (T_0, ∞) with $\lim_{n \rightarrow \infty} t_n = \infty$ and such that

$$\lim_{n \rightarrow \infty} [G(t_n) - F(t_n)] = \limsup_{t \rightarrow \infty} [G(t) - F(t)].$$

In view of (12), there exists a constant M such that

$$(18) \quad G(t_n) - F(t_n) \leq M \quad \text{for all sufficient large } n.$$

It follows from (17) that

$$(19) \quad \lim_{n \rightarrow \infty} G(t_n) = \infty.$$

This and (18) give

$$(20) \quad \lim_{n \rightarrow \infty} F(t_n) = \infty.$$

Then, by (18) and (19),

$$\frac{F(t_n)}{G(t_n)} - 1 \geq \frac{-M}{G(t_n)} > \frac{-1}{2} \quad \text{for } n \text{ large enough.}$$

Thus,

$$\frac{F(t_n)}{G(t_n)} > \frac{1}{2} \quad \text{for } n \text{ large enough.}$$

This and (20) imply that

$$(21) \quad \lim_{n \rightarrow \infty} \frac{F^\alpha(t_n)}{G^{\alpha-1}(t_n)} = \infty.$$

On the other hand, by the Holder's inequality, we have

$$\begin{aligned} F(t_n) &= \frac{1}{H(t_n, T_0)} \int_{T_0}^{t_n} |h(t_n, s)| |\omega(s)| ds \\ &\leq \left\{ \frac{\alpha - 1}{H(t_n, T_0)} \int_{T_0}^{t_n} H(t_n, s) k(s) \frac{|\omega(s)|^{\alpha/(\alpha-1)}}{[\rho(s)r(s)]^{1/(\alpha-1)}} ds \right\}^{(\alpha-1)/\alpha} \\ &\quad \times \left\{ \frac{1}{(\alpha - 1)^{\alpha-1} H(t_n, T_0)} \int_{T_0}^{t_n} \frac{\rho(s)r(s) |h(t_n, s)|^\alpha}{(H(t_n, s)k(s))^{\alpha-1}} ds \right\}^{1/\alpha} \\ &\leq \frac{G^{(\alpha-1)/\alpha}(t_n)}{(\alpha - 1)^{(\alpha-1)/\alpha}} \left\{ \frac{1}{H(t_n, T_0)} \int_{T_0}^{t_n} \frac{\rho(s)r(s) |h(t_n, s)|^\alpha}{(H(t_n, s)k(s))^{\alpha-1}} ds \right\}^{1/\alpha}, \end{aligned}$$

and therefore,

$$\begin{aligned} \frac{F^\alpha(t_n)}{G^{\alpha-1}(t_n)} &\leq \frac{1}{(\alpha - 1)^{(\alpha-1)} H(t_n, T_0)} \int_{T_0}^{t_n} \frac{\rho(s)r(s) |h(t_n, s)|^\alpha}{(H(t_n, s)k(s))^{\alpha-1}} ds \\ &\leq \frac{1}{(\alpha - 1)^{(\alpha-1)} \eta H(t_n, t_0)} \int_{t_0}^{t_n} \frac{\rho(s)r(s) |h(t_n, s)|^\alpha}{(H(t_n, s)k(s))^{\alpha-1}} ds, \end{aligned}$$

for all large n . It follows from (21) that

$$(22) \quad \lim_{n \rightarrow \infty} \frac{1}{H(t_n, t_0)} \int_{t_0}^{t_n} \frac{\rho(s)r(s) |h(t_n, s)|^\alpha}{(H(t_n, s)k(s))^{\alpha-1}} ds = \infty,$$

that is,

$$\lim_{t \rightarrow \infty} \frac{1}{H(t, t_0)} \int_{T_0}^t \frac{\rho(s)r(s) |h(t, s)|^\alpha}{(H(t, s)k(s))^{\alpha-1}} ds = \infty,$$

which contradicts (7). Hence, (13) holds. Then, it follows from (10) that

$$\int_{T_0}^t \frac{\Omega_+^{\alpha/(\alpha-1)}(s)}{[k(s)\rho(s)r(s)]^{1/(\alpha-1)}} ds \leq \int_{T_0}^t k(s) \frac{|\omega(s)|^{\alpha/(\alpha-1)}}{[\rho(s)r(s)]^{1/(\alpha-1)}} ds < \infty,$$

which contradicts (8). This completes the proof of Theorem 2. \square

Theorem 3. *Suppose that (1) and (2) hold, and let the functions H, h, ρ and k be the same as in Theorem 1. such that (6) and*

$$(23) \quad \liminf_{t \rightarrow \infty} \frac{1}{H(t, t_0)} \int_{t_0}^t H(t, s)\rho(s)k(s)q(s)ds < \infty,$$

hold. If there exists a function $\Omega \in C([t_0, \infty), \mathbf{R})$ such that (8) hold for every $T \geq t_0$ and

$$(24) \quad \liminf_{t \rightarrow \infty} \frac{1}{H(t, T)} \int_T^t \left[H(t, s)\rho(s)k(s)q(s) - \frac{\gamma\rho(s)r(s) |h(t, s)|^\alpha}{\alpha^\alpha (H(t, s)k(s))^{\alpha-1}} \right] ds \geq \Omega(T),$$

then equation (E_1) is oscillatory.

Proof. Without loss of generality, we may assume that there exists a solution $x(t)$ of equation (E_1) such that $x(t) \neq 0$ on $[T_0, \infty)$ for some sufficiently large $T_0 \geq t_0$. Define $\omega(t)$ as of Theorem 1. As in the proofs of Theorem 1 and 2, we can obtain (4), (5) and (10). From (23) it follow that

$$(25) \quad \begin{aligned} \limsup_{t \rightarrow \infty} |G(t) - F(t)| &\leq k(t_0)\omega(t_0) \\ &\quad - \liminf_{t \rightarrow \infty} \frac{1}{H(t, t_0)} \int_{t_0}^t H(t, s)\rho(s)k(s)q(s)ds \\ &< \infty, \end{aligned}$$

where $F(t)$ and $G(t)$ are defined as in the proof of Theorem 2. By (24) we have

$$\begin{aligned} \Omega(t_0) &\leq \liminf_{t \rightarrow \infty} \frac{1}{H(t, t_0)} \int_{t_0}^t H(t, s)\rho(s)k(s)q(s)ds \\ &\quad - \liminf_{t \rightarrow \infty} \frac{1}{H(t, t_0)} \int_{t_0}^t \frac{\gamma\rho(s)r(s) |h(t, s)|^\alpha}{\alpha^\alpha (H(t, s)k(s))^{\alpha-1}} ds. \end{aligned}$$

This and (23) imply that

$$\liminf_{t \rightarrow \infty} \frac{1}{H(t, t_0)} \int_{t_0}^t \frac{\rho(s)r(s) |h(t, s)|^\alpha}{(H(t, s)k(s))^{\alpha-1}} < \infty,$$

considering a sequence $\{t_n\}_{n=1}^\infty$ in (T_0, ∞) with $\lim_{n \rightarrow \infty} t_n = \infty$ and such that

$$\begin{aligned} & \lim_{n \rightarrow \infty} \frac{1}{H(t_n, t_0)} \int_{t_0}^{t_n} \frac{\rho(s)r(s) |h(t_n, s)|^\alpha}{(H(t_n, s)k(s))^{\alpha-1}} ds \\ (26) \quad & = \liminf_{t \rightarrow \infty} \frac{1}{H(t, t_0)} \int_{t_0}^t \frac{\rho(s)r(s) |h(t, s)|^\alpha}{(H(t, s)k(s))^{\alpha-1}} ds < \infty. \end{aligned}$$

Now, suppose that (14) holds. With the same argument as in Theorem 2, we conclude that (17) is satisfied. By (25), there exists a constant M such that (18) is fulfilled. Then, following the procedure of the proof of Theorem 2, we see that (22) holds, which contradicts (26). This contradiction proves that (26) fails. The remainder of the proof is similar to that of Theorem 2, so we omit the details. This completes the proof of Theorem 3. \square

Theorem 4. *Suppose that (1) and (2) hold, and let the functions H, h, ρ and k be the same as in Theorem 1 such that (6), and*

$$(27) \quad \liminf_{t \rightarrow \infty} \frac{1}{H(t, t_0)} \int_{t_0}^t \frac{\rho(s)r(s) |h(t, s)|^\alpha}{(H(t, s)k(s))^{\alpha-1}} ds < \infty,$$

hold. If there exists a function $\Omega \in C([t_0, \infty), \mathbf{R})$ such that (8) and (24) hold for every $T \geq t_0$, then equation (E_1) is oscillatory.

Remark 1. If $p(t) \equiv 0$, and $\psi(x) \equiv 1$, then the above Theorems 1, 2 and 4 extend and improve Theorems 1, 2 and 3 of Manojlovic [12], and Theorems 1 – 3 reduce to Theorems 1 – 3 of Wang [14], respectively.

Remark 2. If $p(t) \equiv 0$, then Theorems 1 – 4, extend and improve Theorems 1 – 4 of Wu et al., [16].

Example 1. Consider the differential equation

$$\frac{d}{dt} \left(t^{-\frac{3}{2}}(1 + e^{-|x(t)|}) \frac{dx}{dt} \right) + t^{-5/2}x(t) = 0 \text{ for } t \geq t_0 > 0.$$

We note that

$$\alpha = 2, \quad r(t) = t^{-\frac{3}{2}}, \quad q(t) = t^{-5/2} \quad \text{and} \quad \psi(x) = 1 + e^{-|x(t)|}.$$

let

$$\rho(s) = s, \quad k(s) = s \quad \text{and} \quad H(t, s) = (t - s)^2.$$

Then

$$\begin{aligned} & \limsup_{t \rightarrow \infty} \frac{1}{H(t, t_0)} \int_{t_0}^t \left[H(t, s) \rho(s) k(s) q(s) - \frac{\gamma \rho(s) r(s) |h(t, s)|^\alpha}{\alpha^\alpha (H(t, s) k(s))^{\alpha-1}} \right] ds \\ &= \limsup_{t \rightarrow \infty} \frac{1}{(t - t_0)^2} \int_{t_0}^t \left(t^2 s^{-\frac{1}{2}} - 2ts^{\frac{1}{2}} + s^{\frac{3}{2}} - 8s^{-\frac{1}{2}} + 8s^{-\frac{1}{2}}t - s^{-\frac{3}{2}}t^2 \right) ds = \infty. \end{aligned}$$

Hence, this equation is oscillatory by Theorem 1. However, none of the results of [12], [14] and [16] are applicable for this equation.

Example 2. Consider the differential equation

$$\frac{d}{dt} \left(\frac{2 + \cos^2(\ln t)}{1 + 3 \cos^2(\ln t)} \frac{1 + 3x^2(t)}{2 + x^2(t)} \frac{d}{dt} x(t) \right) + \frac{1}{t} \frac{d}{dt} x(t) + \frac{1}{t^2} x(t) = 0,$$

for $t \geq t_0 = 1$. We note that

$$\alpha = 2, \quad r(t) = \frac{2 + \cos^2(\ln t)}{1 + 3 \cos^2(\ln t)}, \quad q(t) = \frac{1}{t^2}, \quad p(t) = \frac{1}{t},$$

and

$$0 < \psi(x) = \frac{1 + 3x^2}{2 + x^2} \leq 3.$$

If we take

$$\rho(s) = 1, \quad k(s) = s \quad \text{and} \quad H(t, s) = (t - s)^2,$$

then $h(t, s) = 2s(t - s)$ and

$$\begin{aligned} & \limsup_{t \rightarrow \infty} \frac{1}{H(t, t_0)} \int_{t_0}^t \left[H(t, s) \rho(s) k(s) q(s) - \frac{\gamma \rho(s) r(s) |h(t, s)|^\alpha}{\alpha^\alpha (H(t, s) k(s))^{\alpha-1}} \right] ds \\ &= \limsup_{t \rightarrow \infty} \frac{1}{(t - 1)^2} \int_1^t \left[(t - s)^2 \frac{1}{s} - 3s \left(\frac{2 + \sin^2(\ln s)}{1 + 3 \sin^2(\ln s)} \right) \right] ds \\ &\geq \limsup_{t \rightarrow \infty} \frac{1}{(t - 1)^2} \int_1^t \left[(t - s)^2 \frac{1}{s} - 6s \right] ds \end{aligned}$$

$$= \limsup_{t \rightarrow \infty} \frac{1}{(t-1)^2} \left[t^2 \ln t - \frac{9}{2}t^2 + 2t + \frac{5}{2} \right] = \infty.$$

Hence by Theorem 1 this equation is oscillatory. One such solution of this equation is $x(t) = \cos(\ln t)$.

3. Oscillation criteria for (E_2) .

Theorem 5. *Suppose that (1) and*

$$(28) \quad \frac{\frac{d}{dx}g(x)}{(\psi(x)|g(x)^{\alpha-2}|)^{1/(\alpha-1)}} \geq \delta > 0 \quad \text{for } x \neq 0,$$

hold, and let the functions H , h , ρ and k be the same as in Theorem 1. Then equation (E_2) is oscillatory if

$$\limsup_{t \rightarrow \infty} \frac{1}{H(t, t_0)} \int_{t_0}^t \left[H(t, s)\rho(s)k(s)q(s) - \frac{\beta^{1-\alpha}\rho(s)r(s)|h(t, s)|^\alpha}{\alpha^\alpha (H(t, s)k(s))^{\alpha-1}} \right] ds = \infty,$$

where $\beta = \frac{\delta}{\alpha-1}$.

Proof. Let $x(t)$ a nonoscillatory solution of equation (E_2) . Without loss of generality, we may assume that $x(t) \neq 0$ on $[T_0, \infty)$ for some sufficiently large $T_0 \geq t_0$. Define $\omega(t)$ as

$$\omega(t) = \rho(t) \frac{r(t)\psi(x) \left| \frac{dx}{dt} \right|^{\alpha-2} \frac{dx}{dt}}{g(x)} \quad \text{for } t \geq t_0.$$

Thus,

$$\begin{aligned} \frac{d}{dt}\omega(t) &= \frac{d}{dt}\rho(t) \frac{r(t)\psi(x) \left| \frac{dx}{dt} \right|^{\alpha-2} \frac{dx}{dt}}{g(x)} + \rho(t) \frac{\frac{d}{dt} \left(r(t)\psi(x) \left| \frac{dx}{dt} \right|^{\alpha-2} \frac{dx}{dt} \right)}{g(x)} \\ &\quad - \frac{\frac{d}{dx}g(x)}{(\psi(x)|g(x)^{\alpha-2}|)^{1/(\alpha-1)}} \frac{|\omega(t)|^{\frac{\alpha}{\alpha-1}}}{[\rho(t)r(t)]^{1/(\alpha-1)}}. \end{aligned}$$

This and equation (E_2) imply

$$\frac{d}{dt}\omega(t) \leq \frac{\frac{d}{dt}\rho(t)}{\rho(t)}\omega(t) - \rho(t)[q(t) + p(t)\varphi(1, \frac{\omega(t)}{\rho(t)})] - \frac{\frac{d}{dx}g(x)}{(\psi(x)|g(x)^{\alpha-2}|)^{1/(\alpha-1)}} \frac{|\omega(t)|^{\frac{\alpha}{\alpha-1}}}{[\rho(t)r(t)]^{1/(\alpha-1)}}.$$

From (1) and (28) we have

$$\frac{d}{dt}\omega(t) \leq \frac{\frac{d}{dt}\rho(t)}{\rho(t)}\omega(t) - \rho(t)q(t) - p(t)\omega(t) - \frac{\delta}{[\rho(t)r(t)]^{1/(\alpha-1)}} |\omega(t)|^{\frac{\alpha}{\alpha-1}}.$$

Multiply the above inequality by $H(t, s)k(s)$ and integrate from T to t we obtain

$$\int_T^t H(t, s)k(s)\rho(s)q(s)ds \leq \int_T^t H(t, s)k(s) \left(\frac{\frac{d}{ds}\rho(s)}{\rho(s)} - p(s) \right) \omega(s)ds - \int_T^t H(t, s)k(s) \frac{d}{ds}\omega(s)ds - \delta \int_T^t H(t, s)k(s)[\gamma\rho(s)r(s)]^{\frac{-1}{\alpha-1}} |\omega(s)|^{\frac{\alpha}{\alpha-1}} ds.$$

Since

$$- \int_T^t H(t, s)k(s) \frac{d}{ds}\omega(s)ds = H(t, T)k(T)\omega(T) + \int_T^t \frac{\partial}{\partial s}(H(t, s)k(s))\omega(s)ds.$$

The previous inequality becomes

$$(29) \quad \int_T^t H(t, s)k(s)\rho(s)q(s)ds \leq H(t, T)k(T)\omega(T) + \int_T^t |h(t, s)\omega(s)| ds - (\alpha - 1) \int_T^t \frac{\beta H(t, s)k(s) |\omega(s)|^{(\alpha-1)/\alpha}}{[\rho(s)r(s)]^{1/(\alpha-1)}} ds.$$

Define

$$X = \frac{1}{\alpha} \left[\beta^{(1-\alpha)/\alpha} [H(t, s)k(s)]^{(1-\alpha)/\alpha} [\rho(s)r(s)]^{1/\alpha} |h(t, s)| \right],$$

$$Y = \left(\beta^{(\alpha-1)/\alpha} [H(t, s)k(s)]^{(\alpha-1)/\alpha} [\rho(s)r(s)]^{-1/\alpha} |\omega(s)| \right)^{1/(\alpha-1)}.$$

Then use the lemma 1, we have

$$|h(t, s)\omega(s)| - (\alpha - 1) \frac{\beta H(t, s)k(s) |\omega(s)|^{\frac{\alpha}{\alpha-1}}}{[\rho(s)r(s)]^{1/(\alpha-1)}} \leq \frac{\beta^{1-\alpha} \rho(s)r(s) |h(t, s)|^\alpha}{\alpha^\alpha (H(t, s)k(s))^{\alpha-1}}.$$

From (29) we have

$$\int_T^t H(t, s)k(s)\rho(s)q(s)ds \leq H(t, T)k(T)\omega(T) + \int_T^t \frac{\beta^{1-\alpha} \rho(s)r(s) |h(t, s)|^\alpha}{\alpha^\alpha (H(t, s)k(s))^{\alpha-1}} ds,$$

or

$$\int_T^t \left[H(t, s)k(s)\rho(s)q(s) - \frac{\beta^{1-\alpha} \rho(s)r(s) |h(t, s)|^\alpha}{\alpha^\alpha (H(t, s)k(s))^{\alpha-1}} \right] ds \leq H(t, T)k(T)\omega(T).$$

The remainder of the proof proceeds as in the proof of Theorem 1. The proof is complete. \square

Following the procedure of the proof of Theorem 2 and 3, we can also prove the following theorems.

Theorem 6. *Suppose that (1) and (28) hold, and let the functions H , h , ρ and k be the same as in Theorem 1. If there exist a functions $\Omega \in C([t_0, \infty), \mathbf{R})$ such that*

$$(30) \quad \liminf_{t \rightarrow \infty} \frac{1}{H(t, t_0)} \int_{t_0}^t \frac{\rho(s)r(s) |h(t, s)|^\alpha}{(H(t, s)k(s))^{\alpha-1}} ds < \infty,$$

and that for every $T \geq t_0$,

$$(31) \quad \liminf_{t \rightarrow \infty} \frac{1}{H(t, T)} \int_T^t \left[H(t, s)\rho(s)k(s)q(s) - \frac{\beta^{1-\alpha} \rho(s)r(s) |h(t, s)|^\alpha}{\alpha^\alpha (H(t, s)k(s))^{\alpha-1}} \right] ds \geq \Omega(T),$$

and (8) hold, then every solution of (E_2) is oscillatory.

Theorem 7. *Suppose that (1) and (28) hold, and let the functions H , h , ρ and k be the same as in Theorem 1. If there exist a function $\Omega \in C([t_0, \infty), \mathbf{R})$ such that*

$$(32) \quad \liminf_{t \rightarrow \infty} \frac{1}{H(t, T)} \int_T^t H(t, s)\rho(s)k(s)q(s)ds < \infty,$$

and that (8) and (31) hold, then every solution of (E_2) is oscillatory.

Theorem 8. Suppose that (1) and (28) hold, and let the functions H , h , ρ and k be the same as in Theorem 1. If there exist a function $\Omega \in C([t_0, \infty), \mathbf{R})$ such that

$$(33) \quad \limsup_{t \rightarrow \infty} \frac{1}{H(t, t_0)} \int_{t_0}^t \frac{\rho(s)r(s)|h(t, s)|^\alpha}{(H(t, s)k(s))^{\alpha-1}} ds < \infty,$$

and that for every $T \geq t_0$,

$$(34) \quad \limsup_{t \rightarrow \infty} \frac{1}{H(t, T)} \int_T^t \left[H(t, s)\rho(s)k(s)q(s) - \frac{\beta^{1-\alpha}\rho(s)r(s)|h(t, s)|^\alpha}{\alpha^\alpha (H(t, s)k(s))^{\alpha-1}} \right] ds \geq \Omega(T),$$

and (8) hold, then every solution of (E_2) is oscillatory.

Remark 3. If $p(t) \equiv 0$, then Theorem 5, 6 and 8 extend and improve Theorem 1, 3 and 2 of Manojlovic [13].

Example 3. Consider the differential equation

$$\frac{d}{dt} \left(t^{-1}x^2(t) \frac{dx(t)}{dt} \right) + t^{-2}x^3(t) = 0 \quad \text{for } t \geq t_0 > 0.$$

We note that

$$\alpha = 2 \quad \text{and} \quad \frac{\frac{d}{dx}g(x)}{\psi(x)} = 3.$$

Let

$$\rho(s) = 1, \quad k(s) = s \quad \text{and} \quad H(t, s) = (t - s)^2.$$

Then

$$\begin{aligned} & \limsup_{t \rightarrow \infty} \frac{1}{H(t, t_0)} \int_{t_0}^t \left[H(t, s)\rho(s)k(s)q(s) - \frac{\beta^{1-\alpha}\rho(s)r(s)|h(t, s)|^\alpha}{\alpha^\alpha (H(t, s)k(s))^{\alpha-1}} \right] ds \\ &= \limsup_{t \rightarrow \infty} \frac{1}{(t - t_0)^2} \int_{t_0}^t \left\{ \frac{t^2}{s} - 2t + s - \frac{t^2}{12s} + \frac{t}{2s} - \frac{3}{4} \right\} ds = \infty. \end{aligned}$$

Hence, this equation is oscillatory by Theorem 5. However the result of Manojlovic [13] do not apply to this equation.

Example 4. Consider the differential equation

$$\frac{d}{dt} \left(\frac{2 + \sin^2(\ln t)}{1 + 3 \sin^2(\ln t)} \frac{1 + 3x^2(t)}{2 + x^2(t)} \frac{dx}{dt}(t) \right) + \frac{1}{t} \frac{dx}{dt} + \frac{(x(t) + x^3(t))}{t^2(1 + 3 \sin^2(\ln t))} = 0,$$

for $t \geq t_0 = 1$. We note that

$$\frac{\frac{d}{dx}g(x)}{\psi(x)} = 2 + x^2 \geq 2 = \delta.$$

If we take

$$\rho(s) = 1, \quad k(s) = s \quad \text{and} \quad H(t, s) = (t - s)^2,$$

then $h(t, s) = 2s(t - s)$ and

$$\begin{aligned} & \limsup_{t \rightarrow \infty} \frac{1}{H(t, t_0)} \int_{t_0}^t \left[H(t, s) \rho(s) k(s) q(s) - \frac{\beta^{1-\alpha} \rho(s) r(s) |h(t, s)|^\alpha}{\alpha^\alpha (H(t, s) k(s))^{\alpha-1}} \right] ds \\ &= \limsup_{t \rightarrow \infty} \frac{1}{(t-1)^2} \int_1^t \left[\left(\frac{(t-s)^2}{s(1+3\sin^2(\ln s))} \right) - \frac{s}{2} \left(\frac{2+\sin^2(\ln s)}{1+3\sin^2(\ln s)} \right) \right] ds \\ &\geq \limsup_{t \rightarrow \infty} \frac{1}{(t-1)^2} \int_1^t \left[\left(\frac{(t-s)^2}{4s} \right) - s \right] ds \\ &= \limsup_{t \rightarrow \infty} \frac{1}{(t-1)^2} \left[\frac{1}{4} t^2 \ln t - \frac{7}{8} t^2 + \frac{1}{2} t + \frac{3}{8} \right] = \infty. \end{aligned}$$

Hence by Theorem 1 this equation is oscillatory. One such solution of this equation is $x(t) = \sin(\ln t)$.

REFERENCES

- [1] B. AYANLAR, A. TIRYAKI. Oscillation theorems for nonlinear second-Order differential equations. *Comput. Math. Appl.* **44** (2002), 529–538.
- [2] H. B. HSU, C. C. YEH. Oscillation theorems for second order half-linear differential equations. *Appl. Math. Lett.* **9** (1996), 71–77.

- [3] G. H. HARDY, J. E. LITTLEWOOD, G. POLYA. Inequalities, 2nd edition, Cambridge University Press, Cambridge, 1988.
- [4] T. KUSANO, Y. NAITO. Oscillation and nonoscillation criteria for second order quasilinear differential equations. *Acta Math. Hungar.* **76**, 1–2 (1997), 81–99.
- [5] T. KUSANO, N. YOSHIDA. Nonoscillation theorems for a class of quasilinear differential equations of second order. *J. Math. Anal. Appl.* **189** (1995), 115–127.
- [6] H. J. LI, C. C. YEH. An integral criterion for oscillation of nonlinear differential equations. *Math. Japon.* **41** (1995), 185–188.
- [7] H. J. LI, C. C. YEH. Oscillation criteria for nonlinear differential equations. *Houston J. Math.* **21** (1995), 801–811.
- [8] H. J. LI, C. C. YEH. Nonoscillation criteria for second order half-linear differential equations. *Appl. Math. Lett.* **8** (1995), 63–70.
- [9] H. J. LI, C. C. YEH. Nonoscillation theorems for second order quasilinear differential equations. *Publ. Math. Debrecen* **47**, 3–4 (1995), 271–279.
- [10] H. J. LI, C. C. YEH. Oscillation of half-linear second order differential equations. *Hiroshima Math. J.* **25** (1995), 585–594.
- [11] W. C. LIAN, C. C. YEH, H. J. LI. The distance between zeros of an oscillatory solution to a half-linear differential equations. *Comput. Math. Appl.* **29** (1995), 39–43.
- [12] J. V. MANOJLOVIC. Oscillation criteria for second order half-linear differential equations. *Math. Comput. Modelling* **30** (1999), 109–119.
- [13] J. V. MANOJLOVIC. Oscillation theorems for nonlinear differential equations of second order. *Electron. J. Qual. Theory Differ. Equ.* **1** (2000), 1–21.
- [14] J. WANG. On second order quasilinear oscillation. *Funkcial. Ekvac.* **41** (1998), 25–54.
- [15] P. J. Y. WONG, R. P. AGARWAL. Oscillatory behaviour of solutions of certain second order nonlinear differential equations. *J. Math. Anal. Appl.* **198** (1996), 337–354.

- [16] H. W. WU, Q.R. WANG, Y. T. XU. Oscillation and Asymptotics for non-linear second-order differential equations. *Comput. Math. Appl.* **48** (2004), 61–72.

Department of Mathematics

Faculty of Science

Mansoura University

Mansoura, 35516, Egypt

e-mail: emelabbasy@mans.edu.eg

Received September 20, 2006

Revised July 31, 2007