

**DISCRETE MODELS OF TIME-FRACTIONAL DIFFUSION
IN A POTENTIAL WELL**

R. Gorenflo ¹, E. A. Abdel-Rehim ²

*Dedicated to Acad. Bogoljub Stanković,
on the occasion of his 80-th birthday*

Abstract

By generalization of Ehrenfest's urn model, we obtain discrete approximations to spatially one-dimensional time-fractional diffusion processes with drift towards the origin. These discrete approximations can be interpreted (a) as difference schemes for the relevant time-fractional partial differential equation, (b) as random walk models. The relevant convergence questions as well as the behaviour for time tending to infinity are discussed, and results of numerical case studies are displayed.

See also, <http://www.diss.fu-berlin.de/2004/168/index.html>

Mathematics Subject Classification: 26A33, 45K05, 60J60, 60G50, 65N06, 80-99

Key Words and Phrases: generalization of Ehrenfest's urn model, diffusion processes with memory and central drift in a potential well, difference schemes, random walk models, fractional derivative, stochastic processes

1. Introduction

In recent years there has been growing interest in diffusion in various fields of physics, chemistry, and related sciences. It is well known that

the fundamental solution (or Green function) of the classical diffusion (or heat) equation can be interpreted as a Gaussian normal probability density function in space evolving in time.

Robert Brown in 1827 was as far as we know the first interested in diffusion in fluids. He noticed that small particles suspended in fluids perform peculiarly erratic movements. This phenomenon, which can also be observed in gases, is referred to as Brownian motion. It later became clear that Brownian motion is an outward manifestation of the molecular motion postulated by kinetic theory of matter. In 1905 Albert Einstein was the first to develop a satisfactory theory. Later the theory was made more rigorous and extended by Smoluchowski, Fokker, Planck, Burger, Wiener and others. Einstein considered the case of the free particle that is, a particle on which no forces other than those due to the molecules of the surrounding medium are acting. This motion was modelled by the classical diffusion equation (see [28], [32] and [26])

$$\frac{\partial u(x, t)}{\partial t} = a \frac{\partial^2 u(x, t)}{\partial x^2}, a > 0, x \in \mathbb{R}, t > 0. \quad (1.1)$$

The solution $u(x, t)$ of this equation with the initial condition $u(x, 0) = \delta(x - \xi)$ is well known as the corresponding *Green function* or the *fundamental solution*. It represents the probability density of a particle being at the point x in the time instant t when it initially (at the time $t = 0$) is at the point $x = \xi$:

$$u(x, t) = \frac{1}{2\sqrt{\pi at}} e^{-(x-\xi)^2/(4at)}.$$

Here a is a positive constant depending on the temperature, the friction coefficient, the universal gas constant, and finally on the Avogadro number. The free motion of the particle modelled by Eq.(1.1) has met great interest among mathematicians, physicists and others, and it has found many generalizations. Approximating random walks of the particle have also been studied by many authors, see for example, [5], [15]. As soon as the theory for the free particle was established, many modifications in order to take into consideration the external outside force were devised. Assuming the external outside forces acting towards the origin $x = 0$ and being proportional to the distance of the particle from the origin Smoluchowski [26] has shown that Eq.(1.1) on the right-hand side should be augmented by a drift term:

$$\frac{\partial u(x, t)}{\partial t} = a \frac{\partial^2 u(x, t)}{\partial x^2} + \frac{\partial}{\partial x}(bxu(x, t)), b > 0. \quad (1.2)$$

Again we assume that the initial condition is $u(x, 0) = \delta(x - \xi)$. The solution of this equation is [21], [20], [16]

$$u(x, t) = p(\xi; x, t) = \frac{1}{\sqrt{2\pi\sigma(t)}} e^{-\frac{(x-m(t))^2}{2\sigma^2}} \quad x \in \mathbb{R}, t > 0, \quad (1.3)$$

where $m(t) = \xi e^{-bt}$, $\sigma^2 = \frac{a}{b}(1 - e^{-2bt})$, and $a > 0, b > 0$.

As a more general situation, we consider the equation

$$\frac{\partial u(x, t)}{\partial t} = a \frac{\partial^2 u(x, t)}{\partial x^2} - \frac{\partial}{\partial x}(F(x)u(x, t)), \quad a > 0 \quad (1.4)$$

assuming the force $F(x) = -\frac{dU(x)}{dx}$, $U(x)$ to be defined as a symmetric differentiable potential, increasing for $x \geq 0$, $U(x) = U(-x)$. In this general situation of a *potential well*, the drift is directed towards the origin. We pay special attention to the following examples (with $b > 0$):

- (a) $U(x) = \frac{bx^2}{2}$, the harmonic diffusive oscillator
- (b) $U(x) = \frac{bx^4}{4}$, the quartic diffusive oscillator,
- (c) $U(x) = (\frac{dx^2}{2} + \frac{bx^4}{4})$, $d > 0$, an harmonic diffusive oscillator,
- (d) $U(x) = \frac{bx^{2m+2}}{(2m+2)}$, $m = 1, 2, \dots$, the strongly non-linear diffusive oscillator.

Eq.(1.2) can be interpreted as modelling the diffusion of a particle under the action of the external outside force $F(x) = -bx, b > 0$. We should here mention the work of A. Chechkin et al. [3]. These authors deal with the above forms of the potential for diffusion fractional in space, in which $\frac{\partial^2 u(x, t)}{\partial x^2}$ is replaced by a symmetric fractional (in space) derivative.

In our paper, we consider another important generalization of Eq.(1.2), namely we shall discuss the diffusion fractional in time under various forms of forces. By fractional in time, we mean that the first-order time derivative is replaced by the Caputo derivative of order $\beta \in (0, 1]$. In this case Eq.(1.4) goes over into

$$\frac{\partial^\beta u(x, t)}{\partial t^\beta} = a \frac{\partial^2 u(x, t)}{\partial x^2} - \frac{\partial}{\partial x}(F(x)u(x, t)), \quad 0 < \beta \leq 1, a > 0. \quad (1.5)$$

Here we take $-F(x)$ as a differentiable odd function, positive for $x > 0$. The use of the Caputo derivative (see Gorenflo, Mainardi [9], [8], [10], [11],

and the appendix) allows us to take in a natural way into account an initial condition $u(x, 0) = f(x)$ and then consider the evolution of $u(x, t)$ for $x \in \mathbb{R}$, $t > 0$.

We shall present a discrete method of solving approximately Eq.(1.5) with linear drift $F(x) = -bx$ and, by analogy, give the results for the other types of $F(x)$. We shall devise both explicit and implicit differences schemes for $\beta = 1$ and for $0 < \beta < 1$. Then, we shall discuss questions of convergence, in particular for t tending to infinity, and compare the numerical results with the continuous stationary solution of Eq.(1.5). Finally, approximating random walks will be simulated and interpreted.

Let us return to Eq.(1.2) with initial condition $u(x, 0) = \delta(x - \xi)$ and its solution Eq.(1.3). Denoting by $\langle x(t) \rangle = \int_{-\infty}^{\infty} xu(x, t)dx$ the mean position of a corresponding diffusing particle, we see that

$$\langle x(t) \rangle = m(t) = \xi e^{-bt} . \quad (1.6)$$

Now, not going into the details of constructing a solution formula for the more general Eq.(1.5) with $F(x) = -bx$, $u(x, 0) = \delta(x - \xi)$ we remark that we still can calculate the function $\langle x(t) \rangle$. Multiplying Eq.(1.5) by x and integrating over $x \in \mathbb{R}$, using the natural properties $u(x, t) \rightarrow 0$ and $x^n u(x, t) \rightarrow 0$ for all $n \geq 1$ as $|x| \rightarrow \infty$, we get the initial value problem

$$\frac{d^\beta \langle x(t) \rangle}{dt^\beta} = -b \langle x(t) \rangle, \quad 0 < \beta < 1, \quad \langle x(0) \rangle = \xi ,$$

whose solution is (see [11])

$$\langle x(t) \rangle = \xi E_\beta(-bt^\beta) . \quad (1.7)$$

Here $E_\beta(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(1+n\beta)}$ is the Mittag-Leffler function [4]. It is important to compare the asymptotic behaviour in the cases $0 < \beta < 1$ and $\beta = 1$. For $\beta = 1$ we have Eq.(1.6), whereas for $0 < \beta < 1$ we have a power-law decay [11]

$$E_\beta(-bt^\beta) \approx \frac{\sin(\beta\pi) \Gamma(\beta)}{\pi bt^\beta} , \quad t \rightarrow \infty, b > 0 .$$

2. Relation to the Ehrenfest model

We describe a random walk model approximating the diffusion Eq.(1.2) with drift towards the origin. We discretize space and time by a grid $\{(x_j, t_n) \mid j \in \mathbb{Z}, n \in \mathbb{N}_0\}$ with $x_j = jh, t_n = n\tau$. Here $h > 0$ and $\tau > 0$ are the steps in space and in time, respectively. Treating $u(x, t)$ as density of an extensive quantity (like mass, charge, or probability) we want to approximate the collected quantity $\int_{x_j - \frac{h}{2}}^{x_j + \frac{h}{2}} u(x, t_n) dx$ present in a spatial cell $x_j - \frac{h}{2} < x \leq x_j + \frac{h}{2}$ at the instant $t = t_n$ by a clump $y_j^{(n)}$,

$$y_j^{(n)} \approx \int_{x_j - \frac{h}{2}}^{x_j + \frac{h}{2}} u(x, t_n) dx . \tag{2.1}$$

Let us henceforth consider the extensive quantity as probability. Then we will find by symmetric discretization in space and forward differencing in time an approximating random walk model, having the form of an explicit difference scheme for $n \geq 0$:

$$\frac{y_j^{(n+1)} - y_j^{(n)}}{\tau} = a \frac{y_{j+1}^{(n)} - 2y_j^{(n)} + y_{j-1}^{(n)}}{h^2} + \frac{b}{2h} (x_{j+1}y_{j+1}^{(n)} - x_{j-1}y_{j-1}^{(n)}) , \tag{2.2}$$

which we subject to a scaling relation $0 < \mu = \frac{\tau}{h^2}$, as is natural for diffusion problems and usual in numerical analysis, see [25]. Eq.(2.2) is equivalent to the equation

$$y_j^{(n+1)} = \gamma y_j^{(n)} + \lambda_{j+1} y_{j+1}^{(n)} + \rho_{j-1} y_{j-1}^{(n)} , \tag{2.3}$$

with

$$\gamma = (1 - 2a\mu), \lambda_j = a\mu(1 + \frac{j}{R}), \rho_j = a\mu(1 - \frac{j}{R}) \text{ and } R = \frac{2a}{bh^2} .$$

Our intention that Eq.(2.3) should describe a random walk with sojourn probability $y_j^{(n)}$ of a particle being in point x_j at instant t_n (so that Eq.(2.3) describes the transition from instant t_n to t_{n+1}) requires $\gamma \geq 0$ and all occurring $\lambda_j \geq 0, \rho_j \geq 0$, hence $0 < \mu \leq \frac{1}{2a}$ and the restriction of the index j to the set $(-R, -R + 1, -R + 2, \dots, R - 2, R - 1, R)$ of integers with the convenient stipulation that R should be a natural number. We then take $y_j^{(n)} = 0$ for $|j| \geq R + 1$. We keep this in mind for all our discrete schemes.

An objection could be raised against this model. Namely whereas in Eq.(1.2) $-\infty < x < \infty$, we consider Eq.(2.3) by the restriction $-R \leq j \leq R$ only in grid points $x_j = jh$ with $-Rh \leq x_j \leq Rh$. However, as usual in the theory of difference schemes we let tend $h \rightarrow 0$ and by the scaling $\tau = \mu h^2$ also $\tau \rightarrow 0$. Then $Rh = \frac{2a}{bh} \rightarrow \infty$, and thus in the limit the whole real axis $-\infty < x < \infty$ is covered. In this sense the difference scheme Eq.(2.2) with $|j| \leq R$ is consistent to Eq.(1.2). Now as we have the transition probabilities λ_j , ρ_j , and γ of jumping one step of length h to the left, to the right, or staying in place respectively, being non negative and summing up to 1, i.e., $\lambda_j + \rho_j + \gamma = 1$, we can view our discrete model as a Markov chain with a tridiagonal stochastic matrix

$$P = (p_{i,j})_{-R \leq i \leq R, -R \leq j \leq R} \text{ with } p_{i,j} = 0 \quad \forall |i - j| \geq 2,$$

and $p_{i,i} = \gamma$, $p_{i,i+1} = \rho_i$ and $p_{i,i-1} = \lambda_i$. Then Eq.(2.3) can be written in the condensed form

$$y^{(n+1)} = P^T \cdot y^{(n)}, \quad n \in \mathbb{N}_0, \quad (2.4)$$

with the column vector, $y^{(k)} = \{y_{-R}^{(k)}, y_{-R+1}^{(k)}, \dots, y_R^{(k)}\}^T$, $k \in \mathbb{N}_0$. For the interpretation of $y^{(n)}$ as a vector of probabilities, we require $y^{(0)}$ likewise to be such column vector, namely $y_j^{(0)} \geq 0$ and $\sum_{j=-R}^R y_j^{(0)} = 1$. Then all $y^{(n)}$ for $n = 1, 2, 3, \dots$ are non-negative. Furthermore,

$$\sum_{j=-R}^R y_j^{(n)} = 1 \quad \forall n \in \mathbb{N}_0,$$

as is easily shown by induction. Actually, the evolution of $y^{(n)}$ is that of a Markov chain [5] with possible states $(x_{-R}, x_{-R+1}, \dots, x_{R-1}, x_R)$.

We compare this discrete model with the generalized Ehrenfest model described by Vincze [31]. Vincze considers N balls, numbered from 1 to N , k of them in an urn U_1 , $N - k$ in an urn U_2 . In an urn U_0 there are $N + s$ slips of paper ($s \geq 0$) each of them having probability $(N + s)^{-1}$ of being randomly drawn. N of the slips are numbered from 1 to N , the other s slips are not numbered. We repeat indefinitely the following experiment.

We draw a slip from the urn U_0 . If it carries a number we move the ball which has the same number from the urn (U_1 or U_2) in which it is lying to the other urn (U_2 or U_1). If the slip is not numbered, we leave the ball in its

urn. Then we put the slip back into the urn U_0 . If we record the states as the number of balls in the urn U_1 , then there are three probabilities: $\frac{k}{N+s}$ for the next state to be $k-1$, $\frac{N-k}{N+s}$ for the next state to be $k+1$, and finally $\frac{s}{N+s}$ for the next state to be k again. In the special case ($s=0$) we have the classical Ehrenfest model described by many authors (see [23], [15], [1], [7], [29], and [2]).

In other words, if $x_n^{(s)} = k$ is the number of balls in urn U_1 after n steps, we denote the respective probability by $q_k^{(n)}$ and get the transition probabilities

$$q_{k,k} = P(x_{n+1}^{(s)} = k | x_n^{(s)} = k) = \frac{s}{s+N}, \quad k = 0, 1, \dots, N, \quad ,$$

$$q_{k,k-1} = P(x_{n+1}^{(s)} = k-1 | x_n^{(s)} = k) = \frac{k}{s+N}, \quad k = 1, 2, \dots, N, \quad ,$$

$$q_{k,k+1} = P(x_{n+1}^{(s)} = k+1 | x_n^{(s)} = k) = \frac{N-k}{s+N}, \quad k = 0, 1, 2, \dots, N-1, \quad ,$$

and

$$q_{k,k} + q_{k,k-1} + q_{k,k+1} = 1 \quad \forall k = 0, 1, 2, \dots, N, \quad \quad q_{k,k \pm j} = 0 \quad \forall j \geq 2 \quad .$$

This model is also called the modified Ehrenfest model. In the special case $s=0$, its states can only change by moving one ball from U_1 to U_2 with the probability $q_{k,k-1}$ or one ball from U_2 to U_1 with the probability $q_{k,k+1}$. In the general case $s \geq 0$ the transition probabilities $q_{i,j}$ form a tridiagonal stochastic matrix

$$Q_s = (q_{i,j})_{0 \leq i \leq N, 0 \leq j \leq N},$$

which coincides with the matrix P by the identification $p_{i,j} = q_{i-\frac{N}{2}, j-\frac{N}{2}}$ if N is an even positive integer, $R = \frac{N}{2}$ and $\mu = \frac{N}{2a(s+N)}$. The position $x=0$ corresponds to $N/2$ balls in U_1 , the position $x=Rh$ to N balls in urn U_1 , and the position $x=-Rh$ to 0 balls in urn U_1 . The probability of finding k balls in urn U_1 after $n+1$ steps is, with $q_k^{(n)} = 0$ for $k < 0$ and $k > N$,

$$q_k^{(n+1)} = q_{k-1}^{(n)} \frac{N-k+1}{s+N} + q_k^{(n)} \frac{s}{s+N} + q_{k+1}^{(n)} \frac{k+1}{s+N}, \quad k = 0, 1, \dots, N. \quad (2.5)$$

This equation can be interpreted as a discrete approximation to a diffusion process with central force. It seems that Schrödinger and Kohlrausch were the first to point out the connection between the discrete Ehrenfest model and Brownian motion of an elastically bound particle. Smoluchowski [26]

showed that this model approximates the partial differential Eq.(1.2). This equation describes also the so called Ornstein-Uhlenbeck process [28].

Note the following: In the sense of the urn model s should be an integer, however for Eq.(2.5) to describe a Markov chain, s is only required to be a non-negative real number.

3. Discrete approximation of the time fractional diffusion with central linear drift

In this section we will consider the approximate solution of the Eq.(1.5) with $F(x) = -bx$ obtained by discretizing it by the explicit finite-difference method.

$$D_{\tau * }^{\beta} y_j^{(n+1)} = a \frac{y_{j+1}^{(n)} - 2y_j^{(n)} + y_{j-1}^{(n)}}{h^2} + \frac{b}{2h} (x_{j+1}y_{j+1}^{(n)} - x_{j-1}y_{j-1}^{(n)}), \quad 0 < \beta \leq 1. \quad (3.1)$$

We restrict here $R = \frac{2a}{bh^2}$, as in Section 2, and the index j to the range $(-R, -R+1, -R+2, \dots, R-2, R-1, R)$. Adjusting the spatial step h so that $R \in \mathbb{N}$, we complement Eq.(3.1) by prescribing the non-negative initial values $y_j^{(0)}$ obeying $\sum_{j=-R}^R y_j^{(0)} = 1$, and for convenience, all $y_j^{(n)} = 0$ for $|j| \geq R+1$, $n = 0, 1, 2, \dots$. In Eq.(3.1) the difference operator $D_{\tau * }^{\beta}$ denotes the discretization of the Caputo time derivative (see the Appendix, [12] and [19]). For $\beta = 1$, Eq.(3.1) is reduced to Eq.(1.2). For discretizing the Caputo time derivative, we use, as in [13] a backward Grünwald- Letenikov scheme in time (starting at level $t = t_{n+1}$) which reads

$$D_{\tau * }^{\beta} y_j^{(n+1)} = \sum_{k=0}^{n+1} (-1)^k \binom{\beta}{k} \frac{y_j^{(n+1-k)} - y_j^{(0)}}{\tau^{\beta}}, \quad 0 < \beta \leq 1. \quad (3.2)$$

Observe that $D_{\tau * }^1 y_j^{(n+1)} = \frac{1}{\tau} (y_j^{(n+1)} - y_j^{(n)})$. Note that in case of sufficient smoothness the scheme(3.1) has order of accurecy $O(h^2 + \tau)$. For simplicity and easy writing we take from now on always

$$b = a = 1,$$

hence $R = \frac{2}{h^2}$, and introduce the scaling parameter

$$\mu = \frac{\tau^{\beta}}{h^2}, \quad 0 < \mu \leq \frac{\beta}{2}. \quad (3.3)$$

This specification of a and b only means a special choice of the units of space and time and so there is no restriction of generality. Solving now Eq.(3.2) for $y_j^{(n+1)}$, $-R \leq j \leq R$, gives

$$y_j^{(n+1)} = \sum_{k=0}^n (-1)^k \binom{\beta}{k} y_j^{(0)} + \sum_{k=1}^n (-1)^{k+1} \binom{\beta}{k} y_j^{(n+1-k)} + y_{j+1}^{(n)} \left[\mu + \frac{\mu h^2}{2} (j+1) \right] - 2\mu y_j^{(n)} + y_{j-1}^{(n)} \left[\mu - \frac{\mu h^2}{2} (j-1) \right]. \quad (3.4)$$

Again $y^{(n+1)}$ represents the probability column vector for where to find the particle at the time instant t_{n+1} . It depends on $y_{j-1}^{(n)}, y_j^{(n)}, y_{j+1}^{(n)}, y_j^{(n-1)}, \dots$, and back to $y_j^{(0)}$. This model, having the form of an explicit difference scheme, can be interpreted as a random walk with memory. Let

$$b_n = \sum_{k=0}^n (-1)^k \binom{\beta}{k}, \quad n = 0, 1, 2, \dots; \quad c_k = (-1)^{k+1} \binom{\beta}{k}, \quad k = 1, 2, \dots$$

Then $b_0 = 1$, and all $c_k \geq 0, b_n \geq 0, \sum_{k=1}^{\infty} c_k = 1$. In the case $0 < \beta < 1$ (see [12], [13]) we have $c_1 = \beta > c_2 > c_3 > \dots \rightarrow 0$, and finally the relation

$$b_n + \sum_{k=1}^n c_k = 1. \quad (3.5)$$

By aid of the coefficients b_n and c_k in Eq.(3.4) the expression

$$\sum_{k=1}^n c_k y_j^{(n+1-k)} + b_n y_j^{(0)}$$

represents the dependence on the past (the memory part) and the expression

$$y_{j+1}^{(n)} \left[\mu + \frac{\mu h^2}{2} (j+1) \right] - 2\mu y_j^{(n)} + y_{j-1}^{(n)} \left[\mu - \frac{\mu h^2}{2} (j-1) \right],$$

represents the diffusion part, the particle going one step to the right or one step to the left or remaining in its position. For $\beta = 1$, all c_k for $k \geq 2$ vanish and we have the natural discretization of Eq.(1.2). In Eq.(3.4), $y_j^{(n+1)}$ represents the probability of finding the particle in point x_j at time instant t_{n+1} . Therefore $y_j^{(n+1)}$ must be non-negative because as initial probabilities

all $y_j^{(0)} \geq 0$, and the summation of $y_j^{(n)}$ over the index j at any time t_n must give 1. In other words, we want $s_n = \sum_{j=-R}^R y_j^{(n)} = 1 \quad \forall n \in \mathbb{N}_0$. We prove that this holds by induction. It is true for the initial index $n = 0$. So, as $b_0 = 1$, then

$$y_j^{(1)} = (1 - 2\mu)y_j^{(0)} + y_{j+1}^{(0)}\left[\mu + \frac{\mu h^2}{2}(j+1)\right] + y_{j-1}^{(0)}\left[\mu - \frac{\mu h^2}{2}(j-1)\right]. \quad (3.6)$$

Using $0 < \mu \leq \beta/2$ and taking the stochastic matrix P as in Section 2, we write this equation in matrix-vector form

$$y^{(1)} = P^T \cdot y^{(0)}. \quad (3.7)$$

Now, summing both sides of Eq.(3.6) over j from $-R$ to R , we get

$$s_1 = (1 - 2\mu)s_0 + \mu s_0 + \sum_j \frac{j+1}{R} y_{j+1}^{(0)} + \mu s_0 - \sum_j \frac{j-1}{R} y_{j-1}^{(0)} = s_0.$$

For $n = 1$:

$$y_j^{(2)} = b_1 y_j^{(0)} + (c_1 - 2\mu)y_j^{(1)} + y_{j+1}^{(1)}\left[\mu + \frac{\mu h^2}{2}(j+1)\right] + y_{j-1}^{(1)}\left[\mu - \frac{\mu h^2}{2}(j-1)\right].$$

Here the dependence on the past appears and by the summation we get

$$s_2 = b_1 s_0 + (c_1 - 2\mu)s_1 + \mu s_1 + \sum_j \frac{j+1}{R} y_{j+1}^{(1)} + \mu s_1 - \sum_j \frac{j-1}{R} y_{j-1}^{(1)}.$$

Because of $b_1 + c_1 = 1$, $s_1 = s_0$ we obtain $s_2 = s_0$. Now we assume $s_n = 1$, $n \geq 2$, and to prove that $s_{n+1} = 1$, we write Eq.(3.4) in the form

$$y_j^{(n+1)} = b_n y_j^{(0)} + \sum_{k=1}^n c_k y_j^{(n+1-k)} - 2\mu y_j^{(n)} + \mu \left(1 + \frac{j+1}{R}\right) y_{j+1}^{(n)} + \mu \left(1 - \frac{j-1}{R}\right) y_{j-1}^{(n)}. \quad (3.8)$$

Summing here over j and using Eq.(3.5), we get $s_{n+1} = 1$. So far, we have proved that our difference scheme is conservative and non-negative with respect to its dependence on the past. In fact, $y_j^{(n+1)}$ is a linear combination of all $y_j^{(k)}$ with $-R \leq j \leq R$, $0 \leq k \leq n$, with non-negative coefficients whose sum is 1.

3.1. The solution of the explicit difference scheme

Let us now, in preparation of the next section, treat Eq.(3.4) in matrix-vector notation. We proceed in two steps, considering the step from n to $n + 1$, separately for $n = 0$ and $n \geq 1$. For $n = 0$, transposing the matrix Eq.(3.7) and adopting the notation

$$(y^{(n)})^T = z^{(n)} \quad \forall n \in \mathbb{N} \quad , z^{(n)} = (z_{-R}^{(n)}, z_{-R+1}^{(n)}, \dots, z_R^{(n)}) \quad ,$$

we find

$$z^{(1)} = z^{(0)}.P \quad . \tag{3.9}$$

Here P is the stochastic matrix whose elements are defined in Section 2 whose rows sum to 1 and $z^{(n)}$ is a row vector. It is convenient to write P in the form

$$P = I + \mu H \quad ,$$

with I a unit matrix and H a matrix whose rows sum to zero:

$$H = \begin{pmatrix} -2 & 2 & 0 & \dots & \dots & \dots & \dots & 0 \\ \frac{1}{R} & -2 & (2 - \frac{1}{R}) & \dots & \dots & \dots & \dots & 0 \\ 0 & \frac{2}{R} & -2 & (2 - \frac{2}{R}) & \dots & \dots & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & 0 & (2 - \frac{2}{R}) & -2 & \frac{2}{R} & 0 \\ \dots & \dots & \dots & \dots & 0 & (2 - \frac{1}{R}) & -2 & \frac{1}{R} \\ \dots & \dots & \dots & \dots & \dots & 0 & 2 & -2 \end{pmatrix} \quad .$$

For $n \geq 1$, we define the matrix $Q = c_1 I + \mu H$, and write equation(3.4) as

$$z^{(n+1)} = b_n z^{(0)} + \sum_{k=2}^n c_k z^{(n+1-k)} + z^{(n)}.Q \quad . \tag{3.10}$$

Observe that the matrix Q is stochastic only if $c_1 = 1$ which is equivalent to $\beta = 1$. The restriction for μ in Eq. (3.3) implies that Q is a non-negative matrix.

4. The implicit scheme (Θ -method)

The idea of the θ - method (also known as the weighted method) is to replace $y_j^{(n)}$ in the right hand side of Eq.(3.1) by $(\theta y_j^{(n+1)} + (1 - \theta)y_j^{(n)})$ where $0 < \theta \leq 1$. With $\theta = 0$, we have the explicit scheme. Again, we

shall discuss separately the cases $n = 0$ and $n \geq 1$. For convenience, we set $\bar{\theta} = (1 - \theta)$. For $n = 0$, we rewrite Eq.(3.9) in the form

$$z^{(1)}.(I - \mu\theta H) = z^{(0)}.(I + \mu\bar{\theta}H) . \quad (4.1)$$

Then

$$z^{(1)} = z^{(0)}.(I + \mu\bar{\theta}H).(I - \mu\theta H)^{-1} . \quad (4.2)$$

We want $P_\theta = (I + \mu\bar{\theta}H).(I - \mu\theta H)^{-1}$ to be a stochastic matrix. This wish leads us to conditions for μ depending on θ . First, P_θ must be non-negative. Obviously, $(I - \mu\theta H)$ is a strictly diagonally dominant M-matrix, hence its inverse is non-negative. $(I + \mu\bar{\theta}H)$ is non-negative if $(1 - 2\mu\bar{\theta}) \geq 0$. So we arrive, with $r = 1$ (below to be modified to $r = \beta$), at the conditions [14], [30]

$$0 < \mu \leq r/2 \text{ if } \theta = 0, 0 < \mu \leq \frac{r}{2(1-\theta)} \text{ if } 0 < \theta < 1, 0 < \mu < \infty \text{ if } \theta = 1 .$$

Second, the rows of P_θ must sum to 1. So, we require $P_\theta.\eta = \eta$, with the column vector η all of whose $2R + 1$ components have the value 1, i.e., $\eta = (1, 1, \dots, 1)^T$. Since the rows of H are all summing to zero ($H.\eta = 0$), then $(I - \mu\theta H).\eta = \eta$ and $(I + \mu\bar{\theta}H).\eta = \eta$. Therefore

$$\eta = (I + \mu\bar{\theta}H).\eta = (I + \mu\bar{\theta}H).(I - \mu\theta H)^{-1}.\eta = P_\theta.\eta .$$

Now, we shall prove $\sum_j y_j^{(n)} = 1 \forall n$ which is equivalent to $\sum_j z_j^{(n)} = 1 \forall n$.

To this purpose, we use the simple rule: $w = (w_1, w_2, \dots)^T$ is a probability column vector iff $w_i \geq 0$, $i = 1, 2, \dots$, and $w^T.\eta = 1$.

Now, we know that $(I - \mu\theta H)^{-1}$ and P_θ are stochastic matrices. Hence $z^{(0)}.\eta = 1$ and Eq.(4.2) implies $z^{(1)}.\eta = z^{(0)}.(P_\theta.\eta) = z^{(0)}.\eta = 1$, so the desired relation is true for $n = 0$. For $n \geq 1$, let us rewrite Eq.(3.10) as

$$z^{(n+1)} = b_n z^{(0)} + \sum_{k=1}^n c_k z^{(n+1-k)} + \mu z^{(n)}.H . \quad (4.3)$$

Replacing $z^{(n)}$ in $\mu z^{(n)}.H$ by $\theta z^{(n+1)} + \bar{\theta} z^{(n)}$ and solving for $z^{(n+1)}$ gives

$$z^{(n+1)} = \left[b_n z^{(0)} + \sum_{k=1}^n c_k z^{(n+1-k)} + \mu \bar{\theta} z^{(n)}.H \right] . [I - \mu\theta H]^{-1} . \quad (4.4)$$

Here $(I - \mu\theta H)^{-1}$ is non-negative matrix like all b_n and c_k . Consideration of the index $k = 1$ in Eq. (4.4) however, forces us to sharpen our restriction on μ by taking $r = \beta = c_1$. Hence, assuming $z^{(n)}.\eta = 1$, multiplying Eq.(4.4) by η , and using the important Eq.(3.5), we get $z^{(n+1)}.\eta = 1$. Thus, $z^{(n+1)}$ is a probability row vector. So, $z^{(n)}.\eta = 1$ for all n . As a matter of fact, the implicit scheme allows us to predict the future faster than the explicit scheme, because μ and τ depend on each other by the scaling relation (3.3) and the bound for μ is increasing (up to ∞) if θ is increased from 0 to 1.

5. The convergence to the stationary solution of the model

Vincze, Fritz et al. and Kac (see [31], [6] and [15]) showed that the elements of the iterated stochastic matrix Q_s^n of the generalized discrete Ehrenfest model (Urn model) converge to the binomial distribution at $n \rightarrow \infty$. This means

$$\lim_{n \rightarrow \infty} Q_s^n = \begin{pmatrix} p_0 & p_1 & \dots & p_N \\ \vdots & \vdots & & \vdots \\ p_0 & p_1 & \dots & p_N \end{pmatrix}$$

with

$$p_k = 2^{-N} \binom{N}{k}, \quad k = 0, 1, \dots, N,$$

N being the total number of balls. $p = \{p_0, \dots, p_N\}$ is a probability vector which represents the stationary distribution of the Markov chain whose matrix is Q_s .

Both the probability of finding k balls in the urn U_1 after $n+1$ steps, see Eq.(2.5), and the probability of finding the particle at the point x_j at the time instant t_{n+1} , see Eq.(2.3), are interpreted as discrete approximation to a diffusion with central linear force. By taking the limit as $Rh \rightarrow \infty$ in Eq.(2.3) we have consistency to the partial differential Eq.(1.2) [26]. Since the stochastic matrix P representing the random walk approach for Eq. (2.3) and the stochastic matrix Q_s representing the random walk approach for equation (2.5) are related to each other (see Section 2), the matrix P^n has an analogous limit as $n \rightarrow \infty$. So far, to show the behaviour of this model represented by Eq.(1.2) as $t \rightarrow \infty$ (i.e., for $\beta = 1$), we form a vector \bar{y} with components

$$\bar{y}_j = 2^{-2R} \binom{2R}{j+R}, \quad -R \leq j \leq R, \tag{5.1}$$

and a sequence of numbers $q = \{q(t_1), q(t_2), \dots\}$, where $t_1 < t_2 < \dots \rightarrow \infty$. The number $q(t_i)$ is defined as

$$q(t_i) = \sum_{j=-R}^R |y_j(t_i) - \bar{y}_j|, \quad i = 1, 2, \dots \quad (5.2)$$

The iteration index n is calculated from the relation $n\tau = t_i$, $t_i = 1, 2, \dots \rightarrow \infty$, while τ is calculated from the scaling parameter (3.3). The simulation of the vector q shows that it approximates an exponential function

$$q \approx c e^{-at}, \quad (5.3)$$

where a and c are constants and a is called the rate of exponential convergence. Such exponential convergence is a general property of an ergodic Markov chain (see Feller [5]). The numerical results confirm that a tends to 1 as t tends to infinity (see Fig.15).

For estimating the convergence as $0 < \beta < 1$, i.e. for the non Markovian chain and under the action of the other types of forces (b), (c) and (d), defined in Section 1, we apply the more general method for calculating the discrete stationary solution of Eq.(1.5). This is done by omitting the dependence on time t . First, for $0 < \beta < 1$ and $F(x) = -x$, we omit the dependence on the time in Eq.(4.3). To this purpose, we replace all the indices $n + 1, 0$ and $n + 1 - k$ by simply n . Then Eq.(4.3) converges to $z.H = 0$ which is equivalent to $H^T.y = 0$. This equation is valid for both $\beta = 1$ and $0 < \beta < 1$ and H^T has an eigenvector y^* of eigenvalue zero. Now the vector $\bar{y} = cy^*$ with $c = 1 / \sum_{j=-R}^R y_j^*$ is a vector whose elements sum to 1 and are the same as in Eq.(5.1). Again, we construct a sequence of numbers

$$d = \{d(t_1), d(t_2), \dots\}, \quad (5.4)$$

as in Eq.(5.2). The numerical results for the sequence d for $\beta = 1$ show that it behaves like the vector q Eq.(5.3). The simulation for $0 < \beta < 1$ shows that it approximates a power function

$$d \approx c t^{-\omega}, \quad (5.5)$$

where c and ω are also constants and ω is called the rate of algebraic convergence. The numerical results show that ω tends to β faster than in the case $\beta = 1$. The implicit scheme allows us to calculate the vector d very

fast because the number of steps in this case is less than that of the explicit scheme. For estimating the convergence of the model under the action of the cubic function $F(x) = -x^3$ we apply the same method. The only difference is that we replace the transition probabilities ρ_j and λ_j in the matrix P by

$$\rho'_j = \mu(1 - \frac{j^3}{R^3}) , \lambda'_j = \mu(1 + \frac{j^3}{R^3}) , \tag{5.6}$$

with $-R \leq j \leq R$ and $R = (\frac{2}{h^4})^{1/3}$. The probability $\gamma = 1 - 2\mu$ remains unchanged. The elements q_i and d_i are so small, so we plot their logarithms against t . By using the relation (2.1), we find that the vector \bar{y}/h approximates the normalized exact solution $u(x)$ of the stationary equation of the model (i.e., as $t \rightarrow \infty$). To obtain the stationary solution, we replace $u(x, t)$ by $u(x)$ in Eqs.(1.2 and 1.5). This means, we omit the dependence on time t . In Eq.(1.5) the fractional time derivative $\frac{\partial^\beta u(x, t)}{\partial t^\beta}$ is the *Caputo fractional derivative* (see the Appendix). The *Caputo fractional derivative* of a constant is zero. Then Eq.(1.5) as well as Eq.(1.2) tends to the equation

$$\frac{\partial^2 u(x)}{\partial x^2} - \frac{\partial (F(x)u(x))}{\partial x} = 0 , \tag{5.7}$$

as $t \rightarrow \infty$ whose solution is

$$u(x) = C e^{-U(x)} , \tag{5.8}$$

C being the constant of integration. Since $u(x)$ represents a probability density, we can determine the constant C from the normalization condition $\int_{-\infty}^{+\infty} u(x)dx = 1$. Hence, the solution of Eq.(5.7) for a linear force $-x$ is

$$u(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2} , \tag{5.9}$$

and the stationary solution for the cubic force $-x^3$ is

$$u(x) = \frac{\sqrt{2}}{\Gamma(1/4)} e^{-x^4/4} . \tag{5.10}$$

Another question is that of convergence of the explicit or implicit difference schemes (for a fixed time $t > 0$ and the step length $h \rightarrow 0$) to the exact solution of the fractional diffusion equation Eq.(1.4) (or more generally Eq. (1.5)). We postpone the detailed analysis of this question to another paper, saying here only that it can be done by the method of *inverse isotonicity* which has been applied for a second order parabolic boundary problem in [14]. The essential point is that the difference scheme is of positive type and stable in the sense of numerical analysis.

6. Random walk simulation

We now discuss the discrete random walk model of the elastically bound particle (diffusion under the action of the force $-x$, namely the generalized Ehrenfest model). We shall discuss this model separately for $\beta = 1$ and for $0 < \beta < 1$ to see the effect of the memory part. Firstly, we shall discuss the case $\beta = 1$, namely Eq.(1.2) considered as a Markov chain. If we assume that the particle is sitting at the point x_j at the instant t_n , then as time proceeds to the instant t_{n+1} the particle jumps either to the point x_{j-1} with probability λ_j , to the point x_{j+1} with probability ρ_j , or remains in its position with probability γ . The transition probabilities λ_j , ρ_j , and γ are defined in Section 2. The past history before the instant t_n is completely forgotten. The simulation case is the standard one for Markov chains (see Section 8).

In the case $0 < \beta < 1$ we must keep in mind the whole past history of the wandering particle. Its history consists of its positions at the times $t_0 = 0, t_1, \dots$ and up to t_n . This means, the path of the particle is $x(t_0), x(t_1), \dots, x(t_n)$. The initial position of the particle $x(0) = \xi$ may be any grid point mh inside the interval $[-Rh, Rh]$ and $m \in [-R, R]$. As we have done in the previous sections, we distinguish the cases corresponding to $n = 0$ and $n \geq 1$ in the simulation. For $n = 0$, the random walk of the Eq.(3.6) is Markov-like. For $n \geq 1$ we rewrite Eq.(3.8) as

$$y_j^{(n+1)} = \left(1 - \sum_{k=1}^n c_k\right) y_j^{(0)} + c_n y_j^{(1)} + c_{n-1} y_j^{(2)} + \dots + c_2 y_j^{(n-1)} \\ + (c_1 - 2\mu) y_j^{(n)} + \mu \left(1 + \frac{j+1}{R}\right) y_{j+1}^{(n)} + \mu \left(1 - \frac{j-1}{R}\right) y_{j-1}^{(n)}. \quad (6.1)$$

Clearly all the coefficients $c_1, c_2, \dots, c_n, (1 - \sum_{k=1}^n c_k)$ are non-negative. The idea of simulation is analogous to that of [13]. Assuming the particle sitting at the grid point $x_j \in [-Rh, Rh]$ at instant $t_n, n \geq 1$, then its position at the next instant t_{n+1} is obtained as follows. We set $s_k = \sum_{i=1}^k c_i$ for $k = 1, 2, \dots, n$ and generate a uniformly in $[0, 1]$ distributed random number u . Then we test successively into which one of the intervals $[0, s_1), [s_1, s_2), [s_2, s_3), \dots, [s_n, 1)$ u falls. The length of these intervals are respectively c_1, c_2, \dots, c_n and $b_n = 1 - s_n$. We subdivide the first interval $([0, s_1) = [0, c_1))$ into three sub-intervals of lengths λ_j, γ' and ρ_j where

$\gamma' = (c_1 - 2\mu)$. Now if $u \in [0, c_1)$, we move the particle from its position $x(t_n) = x_j$ to the point x_{j-1} , x_j or x_{j+1} depending on whether u is in the subinterval of length λ_j, γ' or ρ_j , respectively. If $u \in [s_{k-1}, s_k)$ with $2 \leq k \leq n$, we move the particle from its position $x(t_n)$ back to its previous position $x(t_{n+1-k})$. In the case $u \in [s_n, 1)$ we move it back to its initial position $x(t_0) = x(0)$. The sketch of transitions is given in Section 8 (Fig. [31 and 32]).

For the actual simulations let us note the following: for a more general force $F(x)$, we must replace the term $\frac{b}{2h} (x_{j+1}y_{j+1}^{(n)} - x_{j-1}y_{j-1}^{(n)})$ in Eqs. [(2.2) and (3.1)] by $\frac{-1}{2h} (F(x_{j+1})y_{j+1}^{(n)} - F(x_{j-1})y_{j-1}^{(n)})$ and take into account all the other resulting changes in discretization.

7. Conclusion

The explicit difference scheme for the classical diffusion equation $\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}$ can as is well known not only be used for approximating the density $u(x, t)$ but allows also re-interpretation and use as a random walk model for approximate simulation of the path of a particle whose sojourn probability density is $u(x, t)$. Analogous two-fold interpretation of explicit and implicit difference schemes is possible for various types of fractional diffusion problems. If fractionality is in the time variable then the difference scheme takes account of the whole past and consequently its stochastic interpretation yields a backward-oriented random walk in which the whole past history of a particle influences its future (in contrast to forward-oriented continuous or discrete time random walk in which after each jump the past is forgotten). We have here worked out the corresponding theory for time-fractional diffusion with symmetric drift towards the spatial origin and have shown its applicability by carrying out numerical case studies, for both possible uses of the difference schemes. The use of the difference schemes as random walk models in the time-fractional case, however, deserves a comment. The particle paths so produced qualitatively look completely different from those produced by models of continuous time random walk. As both types of models approximate the same sojourn probability density (we plan to say more on this matter in a forthcoming paper) we have here an instance of the fact that totally distinct types of random walk can in the appropriate limit describe the time evolution of the same sojourn probability density.

8. Numerical Results

Figures [1-3] correspond to the explicit scheme for $F(x) = -x$ and $y^{(0)} = \{0, \dots, 1, \dots, 0\}$. Figures [4-6] correspond to the explicit scheme for $F(x) = -x$ and $y^{(0)} = \{\frac{1}{2R+1}, \dots, \frac{1}{2R+1}, \dots, \frac{1}{2R+1}\}$. Figures [7-11] correspond to the implicit scheme for $F(x) = -x$ and $y^{(0)} = \{0, \dots, 1, \dots, 0\}$. Figures [12-14] correspond to the implicit scheme for $F(x) = -x$ and $y^{(0)} = \{0, \dots, 0, 1\}$. Figures [15-16] illustrate the convergence as $f(x) = -x$ and $y^{(0)} = \{0, \dots, 1, \dots, 0\}$. In these figures we have plotted $\log d$ against time t . Figures [17,18] show the approximate stationary solution and the approximate solution of the model of linear force.

Figures [19-20] correspond to the explicit scheme for $F(x) = -x^3$ and $y^{(0)} = \{0, \dots, 0, 1\}$. Figures [21-22] correspond to the implicit scheme for $F(x) = -x^3$ and $y^{(0)} = \{0, \dots, 1, \dots, 0\}$. Figures [23-24] illustrate the convergence of the model for $F(x) = -x^3$ and $y^{(0)} = \{0, \dots, 1, \dots, 0\}$. We have plotted $\log d$ against time t . Figures [25,26] show the approximate stationary solution and the stationary analytic solution of the model for the cubic force $F(x) = -x^3$. Figures [27,30] exhibit the simulation of the random walk and its increments for $F(x) = -x$ and $x(0) = 0$ for $\beta = 1$ and $0 < \beta < 1$. In these figures, we have plotted x or Δx against the number of steps n . The results of all these figures are taken for $R = 10$.




Figure 1:




Figure 2:




Figure 3:




Figure 4:




Figure 5:




Figure 6:

Appendix: Important definitions

For the purpose of this paper, we use the Riemann-Liouville fractional derivative of order $\beta > 0$ such that $0 < \beta < 1$ for a function $f(t)$ given in the interval $[0, b]$, $b < \infty$, defined for $t > 0$ by the expressions (see [22], [11], [18] and [19]).

$$(D^\beta f)(t) = \begin{cases} \frac{1}{\Gamma(1-\beta)} \frac{d}{dt} \int_0^t \frac{f(\tau)}{(t-\tau)^\beta} d\tau, & 0 < \beta < 1 \\ \frac{df(t)}{dt}, & \beta = 1 \end{cases} . \quad (A.1)$$




Figure 7:




Figure 8:




Figure 9:




Figure 10:




Figure 11:




Figure 12:




Figure 13:




Figure 14:




Figure 15:




Figure 16:




Figure 17:




Figure 18:




Figure 19:




Figure 20:




Figure 21:




Figure 22:

We refer the readers to the books [18], [19] and [22] for the general theory of fractional derivatives and integrals. The Riemann-Liouville fractional integral of order $\beta > 0$ is defined as

$$J^\beta f(t) = \frac{1}{\Gamma(\beta)} \int_0^t (t - \tau)^{(\beta-1)} f(\tau) d\tau, \quad t > 0, \quad \beta \in \mathbb{R}^+. \quad (A.2)$$

The alternative fractional derivative operator(see [11]) is the Caputo fractional derivative of order $0 < \beta < 1$.

$$(D_*^\beta f)(t) = \begin{cases} \frac{1}{\Gamma(1-\beta)} \int_0^t \frac{\dot{f}(\tau)}{(t-\tau)^\beta} d\tau, & 0 < \beta < 1 \\ \frac{df(t)}{dt}, & \beta = 1 \end{cases}. \quad (A.3)$$




Figure 23:




Figure 24:




Figure 25:




Figure 26:

This definition is more restrictive than (A.1), and it is instructive to mention the relation between the Riemann-Liouville fractional derivative and integral operators and the Caputo fractional derivative $D_*^\beta f(t) = J^{1-\beta} D^1 f(t)$, $0 < \beta < 1$. Whereas, the Riemann-Liouville fractional derivative and integral operators satisfy the equation $D^\beta f(t) = D^1 J^{1-\beta} f(t)$, $0 < \beta \leq 1$ with $D^1 = \frac{d}{dt}$, we have, if $0 < \beta < 1$,

$$(D_*^\beta f)(t) = D^\beta [f(t) - f(0^+)] = D^\beta f(t) - \frac{f(0)t^{-\beta}}{\Gamma(1-\beta)}. \quad (A.4)$$

In Eq. (1.5) we have written $\frac{\partial^\beta}{\partial t^\beta}$ in place of D_*^β for reasons of notational analogy to $\frac{\partial}{\partial t}$. Eq. (A.4) represents the relation between the Riemann-Liouville




Figure 27:




Figure 28:




Figure 29:




Figure 30:

and Caputo fractional derivative and the dependence on the initial value $f(0)$. It is important for solving fractional differential equations. Finally, since $D^\beta t^\mu = \frac{\Gamma(\mu+1)}{\Gamma(\mu+1-\beta)} t^{\mu-\beta}$, $\beta \geq 0$, $\mu \geq 0$, the Riemann-Liouville fractional derivative of a constant is not zero, but the Caputo fractional derivative of a constant is zero.

Let us now give some information on the Grünwald-Letnikov approximation of fractional derivatives. For more details, see [22], [18] and [19]. For a sufficiently smooth function, defined and bounded on the whole real line, we have

$$f^{(\beta)}(x) = \lim_{h \rightarrow 0^+} \frac{(\Delta_h^\beta f)(x)}{h^\beta}, \quad \beta > 0, \quad (A.5)$$

where $(\Delta_h^\beta f)(x) = \sum_{k=0}^{\infty} \binom{\beta}{k} f(x - kh)$. With the binomial coefficients $\binom{\beta}{k}$ this series converges absolutely and uniformly for each $\beta > 0$ and for every bounded function. Eq.(A.5) is called the Grünwald-Letnikov fractional derivative of order $\beta > 0$ on the whole line. Grünwald-Letnikov defined also the fractional difference of order $\beta > 0$ on a finite interval as

$$(\Delta_h^\beta f)(x) = \sum_{k=0}^{\frac{x-a}{h}} (-1)^k \binom{\beta}{k} f(x - kh), h = \frac{x-a}{n}, n \in \mathbb{N}. \quad (A.6)$$

Here $a \leq x \leq b$, and $f(x)$ is defined only in the interval $[a, b]$, $b > a$. Then, $f^{(\beta)}(x) = \lim_{h \rightarrow 0^+} \frac{1}{h^\beta} (\Delta_h^\beta f)(x)$ is used as reason for our discretization.

Acknowledgements: The authors appreciate support from the Research Commission of Free University of Berlin (Project "Convolutions") and partial support by the *INTAS* project 00-0847. The second named author is grateful for the grant provided by the government of *Arab Republic of Egypt*. Finally, both authors thank an anonymous referee for helpful suggestions.

Remark added in proof (October 2005): The following list of References has been compiled in Spring 2003. Due to the passage of time since then it would deserve an update. To avoid further delay we leave it as it is, apologizing for missing citations. But let us quote the thesis of E. A. Abdel-Rehim (July 2004): *Modelling and Simulation of Classical and Non-Classical Diffusion Processes by Random Walks*. Available at <http://www.diss.fu-berlin.de/2004/168/index.html>.

References

- [1] N. H. Bingham, Fluctuation theory for the Ehrenfest urn. *Adv. Appl. Prob.* **23** (1991), 598-611.
- [2] G. Blom, Mean transition times for the Ehrenfest urn model. *Adv. Appl. Prob.* **21** (1989), 479-480.
- [3] A. Chechekin, V. Gonchar, J. Klafter, R. Metzler, L. Tanatarov, Stationary states of non-linear oscillators driven by Lévy noise. *Chemical Physics* **284** (2002), 233-251.
- [4] A. Erdélyi, W. Magnus, F. Oberhettinger and F. G. Tricomi, *Higher Transcendental Functions, Bateman Project*, Vols. **1-3**. McGraw-Hill, New York (1953-1955).




Figure 31: The sketch of the possible jumps as $\beta = 1$




Figure 32: The sketch of the possible jumps as $0 < \beta < 1$

- [5] W. Feller, *An Introduction to Probability Theory and its Applications*, vol. **1** (3rd ed). New York - London - Sydney (1968).
- [6] F.-J. Fritz, B. Huppert, and W. Willems, *Stochastische Matrizen*. Springer-Verlag, Berlin - Heidelberg - New York (1979).
- [7] C. Godrèche and J. M. Luck, Correlation and response in the backgammon model: the Ehrenfest legacy. *J. Phys. A: Math. Gen.* **32** (1999), 6033-6054.
- [8] R. Gorenflo and F. Mainardi, Approximation of Lévy- Feller diffusion by random walk. *J. for Analysis and its Applications (ZAA)* **18** (1999), 231-146.
- [9] R. Gorenflo and F. Mainardi, Random walk models for space-fractional diffusion processes. *Fractional Calculus and Applied Analysis* **1** (1998), 167-191.
- [10] R. Gorenflo and F. Mainardi, Random walk models approximating symmetric space-fractional diffusion processes. In: J. Elschner, I. Gohberg and B. Silbermann (Editors), *Problems in Mathematical Physics*, Birkhäuser Verlag, Basel (2001), 120-145.
- [11] R. Gorenflo and F. Mainardi, Fractional calculus: Integral and differential equations of fractional order. In: A. Carpinteri and F. Mainardi (Editors). *Fractals and Fractional Calculus in Continuum Mechanics*, Springer Verlag, Wien and New York (1997), 223-276. [Reprinted in News 010101, see <http://www.fractalmo.org>]
- [12] R. Gorenflo, F. Mainardi, D. Moretti, and P. Paradisi, Time-fractional diffusion: A discrete random walk approach. *Nonlinear Dynamics* **29** (2002), 129-143.
- [13] R. Gorenflo, F. Mainardi, D. Moretti, G. Pagnini, and P. Paradisi, Discrete random walk models for space-time fractional diffusion. *Chemical Physics* **284** (2002), 521-541.
- [14] R. Gorenflo and Marianne Niedack, Conservative difference schemes for diffusion problems with boundary and interface conditions. *Computing* **25** (1980), 299-316.
- [15] M. Kac, Random walk and the theory of Brownian motion. *Amer. Math. Monthly* **54** (1947), 369-391.
- [16] N. G. Van Kampen, *Stochastic Processes in Physics and Chemistry*. North-Holland Publishing Company, Amsterdam - New York - Oxford (1981).
- [17] R. Metzler, J. Klafter and I. M. Sokolov, Anomalous transport and external fields: continuous time random walks and fractional diffusion equations extended. *Physical Review E* **58**, No 2 (1998), 1621-1633.
- [18] K.S. Miller and B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*. John Wiley and Sons, New York - Chichester - Brisbane - Toronto - Singapore (1993).
- [19] I. Podlubny, *Fractional Differential Equations*. Academic Press, San Diego - Boston - New York - London - Sydney - Tokyo - Toronto (1999).
- [20] N. U. Prabhu, *Stochastic Processes (Basic Theory and its Applications)*. The Macmillan Company, New York; Collier-Macmillan Limited, London (1965).

- [21] H. Risken, *The Fokker-Planck Equation (Methods of Solution and Applications)* (2nd ed). Springer-Verlag, Berlin - Heidelberg (1989).
- [22] S. G. Samko, A. A. Kilbas, and O. I. Marichev, *Fractional Integrals and Derivatives (Theory and Applications)*. Gordon and Breach, New York - London - Paris (1993). Translated from the Russian edition, Minsk (1987).
- [23] K. Samuel and J. Mc Gregor, Ehrenfest urn models. *J. Appl. Prob.* **2** (1965), 352-376.
- [24] W. R. Schneider and W. Wyss, Fractional diffusion and wave equations. *J. Math. Phys.* **30**, No 1 (1989), 134-144.
- [25] G. D. Smith, *Numerical Solution of Partial Differential Equations with Exercises and Worked Solutions*. Oxford University Press, London - New York - Toronto (1965).
- [26] Smoluchowski, Drei Vorträge über Diffusion, Brownsche Molekularbewegung und Koagulation von Kolloidteilchen. *Phys. Zeit.* **17** (1916), 557-571 and 585-599.
- [27] I. M. Sokolov, J. Klafter, and A. Blumen, Fractional kinetics. *Physics Today* **55** (November 2002), 48-54.
- [28] G.E. Uhlenbeck and L. S. Ornstein, On the theory of Brownian motion. *Phys. Rev.* **36** (1930), 823-841.
- [29] K. W. H. Van Beek and A. J. Stam, A variant of the Ehrenfest model. *Adv. Appl. Prob.* **19** (1987), 995-996.
- [30] R. S. Varga, *Matrix Iterative Analysis*. Inc. Englewood Cliffs, New Jersey, Prentice-Hall (1962).
- [31] I. Vincze, Über das Ehrenfestsche Modell der Wärmeübertragung. *Archiv der Mathematik* **XV**, (1964), 394-400.
- [32] M. C. Wang and G. E. Uhlenbeck, On the theory of Brownian motion II, *Rev. Mod. Phys.* **17** (1956), 323-342.

Department of Mathematics and Computer Science

Free University of Berlin

Arnimalle 3

D-14195 Berlin, GERMANY

Received: June 16, 2003

¹ *e-mail: gorenflo@mi.fu-berlin.de , URL: www.fracalmo.org*

² *e-mail: entsarabdelrehim@yahoo.com*