

EVERY n -DIMENSIONAL SEPARABLE METRIC SPACE
CONTAINS A TOTALLY DISCONNECTED $(n - 1)$ -DIMEN-
SIONAL SUBSET WITH NO TRUE QUASI-COMPONENTS*

Vladimir Todorov, Petar Stoev

The quasi-component $Q(x)$ of a point x of a topological space X is by definition the intersection of all open and closed subsets of X , every one of which contains x . If a quasi-component consists of more than one point, it is called a *true quasi-component*. In this note we give a simple construction of (at least) $(n - 1)$ -dimensional totally disconnected subspace Y of a given n -dimensional separable metric space X such that every quasi-component in Y is a single point.

1. Basic concepts and definitions. Let X be a topological space and $x \in X$ be some point of X . The intersection $Q(x)$ of all closed and open (clopen) subsets of X that contain x is called the *quasi-component* of x . The quasi-component may consist of more than one point even if X does not contain a connected subspace different from a point – for example the Knaster-Kuratowski fan [1].

Suppose, further, that X is a separable metric space and $\dim X = n$. Then X may be regarded as a subset of some Euclidean space \mathbb{R}^m for $m \leq 2n + 1$ [2, p. 262].

A topological space X is called totally disconnected if X does not contain a connected subspace different from a point [3].

And so, we suppose below that $X \subset \mathbb{R}^m$ and $\dim X = n$. We call as well that $F \subset X$ is a *separator* in X , if F is a closed subset of X and $X \setminus F$ is not connected.

2. The space Y . Denote by \mathcal{S} the set of all separators of X . It is easy to see that the cardinality $\text{card}\mathcal{S}$ of \mathcal{S} is equal to \mathfrak{c} , the cardinal number of the continuum.

Consider next the set \mathcal{P} of all hyperplanes in \mathbb{R}^m and denote by \mathcal{P}_0 the subset of \mathcal{P} which consists of hyperplanes p with equations

$$p: a_0 + a_1x_1 + a_2x_2 + \cdots + a_mx_m = 0,$$

such that $a_k, k = 0, 1, \dots, m$ are rational numbers. Evidently, \mathcal{P}_0 is a countable set and, hence, the set $\mathcal{P}_1 = \mathcal{P} \setminus \mathcal{P}_0$ has cardinality \mathfrak{c} . After that it is obvious that to every separator $F \in \mathcal{S}$ one can attach a hyperplane $p_F \in \mathcal{P}_1$ so that the intersection $F \cap p_F$ is non empty set and, moreover, it is possible to choose $x_F \in F \cap p_F$ in such a way that $x_F \neq x_G$ for every two different elements F and G in \mathcal{S} (we suppose here that $\dim X > 1$).

*2000 Mathematics Subject Classification: 17C55.

Key words: Totally disconnected n -dimensional space.

Then, the desired set Y is $\{x_F | F \in \mathcal{S}\}$. Clearly, Y is not connected between any pair $x \neq y$ of different points of Y because one can find a hyperplane $p_{xy} \in \mathcal{P}_0$ which separates x and y in \mathbb{R}^m and, evidently, $p_{xy} \cap Y = \emptyset$.

Next, it is easy to see that $\dim Y \geq n - 1$, because Y meets every partition in X . Note that if we add a single point $\{*\}$ to Y , the space $Y^* = Y \cup \{*\}$ remains totally disconnected, which is an answer of a question of G. Dimov

REFERENCES

- [1] B. KNASTER, K. KURATOWSKI. Sur les ensembles et non connexes. *Fundam. Math.*, **2** (1921), 206–255.
- [2] P. ALEXANDROFF, B. PASINKOV. An introduction to the Dimension Theory. Moscow, Nauka, 1973 (in Russian).
- [3] R. ENGELKING. Outline of general topology. North-Holland, Amsterdam, 1968.

V. Todorov, P. Stoev
Department of Mathematics
University of Architecture, Civil Engineering and Geodesy
1, Hr. Smirnenski Blvd
1421 Sofia, Bulgaria
e-mail: vttp@yahoo.com

ВСЯКО n -МЕРНО СЕПАРАБЕЛНО МЕТРИЧНО ПРОСТРАНСТВО СЪДЪРЖА НАПЪЛНО НЕСВЪРЗАНО $(n - 1)$ -МЕРНО ПОДМНОЖЕСТВО С ЕДНОТОЧКОВИ КВАЗИКОМПОНЕНТИ

Владимир Тодоров, Петър Стоев

Тази бележка съдържа елементарна конструкция на множество с указаните в заглавието свойства. Да отбележим в допълнение, че така полученото множество остава напълно несвързано дори и след като се допълни с краен брой елементи.