

Provided for non-commercial research and educational use.
Not for reproduction, distribution or commercial use.

Serdica

Mathematical Journal

Сердика

Математическо списание

The attached copy is furnished for non-commercial research and education use only.
Authors are permitted to post this version of the article to their personal websites or institutional repositories and to share with other researchers in the form of electronic reprints.
Other uses, including reproduction and distribution, or selling or licensing copies, or posting to third party websites are prohibited.

For further information on
Serdica Mathematical Journal
which is the new series of
Serdica Bulgaricae Mathematicae Publicationes
visit the website of the journal <http://www.math.bas.bg/~serdica>
or contact: Editorial Office
Serdica Mathematical Journal
Institute of Mathematics and Informatics
Bulgarian Academy of Sciences
Telephone: (+359-2)9792818, FAX:(+359-2)971-36-49
e-mail: serdica@math.bas.bg

CAYLEY-HAMILTON THEOREM FOR MATRICES OVER AN ARBITRARY RING

Jenő Szigeti*

Communicated by V. Drensky

ABSTRACT. For an $n \times n$ matrix A over an arbitrary unitary ring R , we obtain the following Cayley-Hamilton identity with right matrix coefficients:

$$(\lambda_0 I + C_0) + A(\lambda_1 I + C_1) + \cdots + A^{n-1}(\lambda_{n-1} I + C_{n-1}) + A^n(nI + C_n) = 0,$$

where $\lambda_0 + \lambda_1 x + \cdots + \lambda_{n-1} x^{n-1} + n! x^n$ is the right characteristic polynomial of A in $R[x]$, $I \in M_n(R)$ is the identity matrix and the entries of the $n \times n$ matrices C_i , $0 \leq i \leq n$ are in $[R, R]$. If R is commutative, then

$$C_0 = C_1 = \cdots = C_{n-1} = C_n = 0$$

and our identity gives the $n!$ times scalar multiple of the classical Cayley-Hamilton identity for A .

2000 *Mathematics Subject Classification*: 15A15, 15A24, 15A33, 16S50.

Key words: The preadjoint and the right characteristic polynomial of an $n \times n$ matrix, the commutator subgroup $[R, R]$ of a ring R .

*The author was supported by OTKA of Hungary, grant no. T43034 and K61007.

1. Introduction. The Cayley-Hamilton theorem and the corresponding trace identity play a fundamental role in proving classical results about the polynomial and trace identities of the $n \times n$ matrix algebra $M_n(K)$ over a field K . In case of $\text{char}(K) = 0$, Kemer's pioneering work (see [2], [3]) on the T-ideals of associative algebras revealed the importance of the identities satisfied by the $n \times n$ matrices over the Grassmann (exterior) algebra

$$E = K \langle v_1, v_2, \dots, v_i, \dots \mid v_i v_j + v_j v_i = 0 \text{ for all } 1 \leq i < j \rangle$$

generated by the infinite sequence of the anticommutative indeterminates $(v_i)_{i \geq 1}$. Accordingly, the importance of matrices over non-commutative rings is an evidence in the theory of PI-rings, nevertheless this fact has been obvious for a long time in other branches of algebra (structure theory of semisimple rings, K-theory, quantum matrices, etc.). Thus a Cayley-Hamilton type identity for such matrices seems to be of general interest.

For $n \times n$ matrices over a Lie-nilpotent ring R a Cayley-Hamilton type identity with one sided scalar coefficients (left or right) was found in [9] (see also in [10]), if R satisfies the PI

$$[[[\dots [[x_1, x_2], x_3], \dots], x_m], x_{m+1}] = 0,$$

then the degree of this identity is n^m . Since E is Lie nilpotent with $m = 2$, the above mentioned identity for a matrix $A \in M_n(E)$ is of degree n^2 . In [1] Domokos presented a slightly modified version of this identity in which the coefficients are invariant under the conjugate action of $GL_n(K)$.

In the general case (when R is an arbitrary non-commutative ring) Paré and Schelter proved (see [4]) that any matrix $A \in M_n(R)$ satisfies a monic identity in which the leading term is A^k for some integer $k \leq 2^{2^{n-1}}$ and the other summands are of the form $r_0 A r_1 A r_2 \dots r_{l-1} A r_l$ with $r_0, r_1, \dots, r_l \in R$ and $0 \leq l \leq k - 1$. An explicit monic identity for 2×2 matrices arising from the argument of [4] and a detailed study of the ideal in $R \langle x \rangle = R * k[x]$ consisting of the polynomials which have as a root the generic $n \times n$ matrix $X = [x_{ij}]$ was given by Robson in [8] ($R = k \langle x_{ij} \rangle$ is the free associative algebra over a field k and $R \langle x \rangle = R * k[x]$ is the free associative k -algebra in one more indeterminate x). Further results in this direction can be found in [5], [6] and [7].

The aim of the present paper is to define the right characteristic polynomial

$$p(x) = \lambda_0 + \lambda_1 x + \dots + \lambda_{n-1} x^{n-1} + n! x^n$$

in $R[x]$ of an $n \times n$ matrix $A \in M_n(R)$ and to derive a corresponding identity for A (here R is an arbitrary unitary ring). We obtain a Cayley-Hamilton identity

with right matrix coefficients of the following form:

$$(\lambda_0 I + C_0) + A(\lambda_1 I + C_1) + \cdots + A^{n-1}(\lambda_{n-1} I + C_{n-1}) + A^n(n! I + C_n) = 0,$$

where $I \in M_n(R)$ is the identity matrix and the entries of the $n \times n$ matrices C_i , $0 \leq i \leq n$ are in the additive subgroup $[R, R]$ of R generated by the commutators $[x, y] = xy - yx$ with $x, y \in R$ (a more precise description of the entries in the C_i 's can be deduced from the proof). Note that a similar identity with left matrix coefficients can be obtained analogously. If R is commutative, then

$$C_0 = C_1 = \cdots = C_{n-1} = C_n = 0$$

and our identity gives the $n!$ times scalar multiple of the classical Cayley-Hamilton identity for A .

We shall make extensive use of the results of [9], in order to provide a self contained treatment, we recall all the necessary prerequisites from [9].

2. The characteristic polynomial. Let R be an arbitrary unitary ring, the *preadjoint* of an $n \times n$ matrix

$$A = [a_{ij}], \quad a_{ij} \in R, \quad 1 \leq i, j \leq n$$

is defined as $A^* = [a_{rs}^*] \in M_n(R)$, where

$$a_{rs}^* = \sum_{\tau, \rho} \operatorname{sgn}(\rho) a_{\tau(1)\rho(\tau(1))} \cdots a_{\tau(s-1)\rho(\tau(s-1))} a_{\tau(s+1)\rho(\tau(s+1))} \cdots a_{\tau(n)\rho(\tau(n))}$$

and the sum is taken over all permutations τ of the set $\{1, \dots, s-1, s+1, \dots, n\}$ and ρ of the set $\{1, 2, \dots, n\}$ with $\rho(s) = r$. The *right determinant* of A is the trace of the product matrix AA^* :

$$r \det(A) = \operatorname{tr}(AA^*).$$

Our development is based on the following crucial result of [9].

Theorem 2.1. *The product $AA^* \in M_n(R)$ can be written in the following form:*

$$AA^* = b_{11} I + C,$$

where b_{11} is the $(1, 1)$ entry in $AA^* = [b_{ij}]$ and $C = [c_{ij}]$ is an $n \times n$ matrix with $c_{11} = 0$ and each c_{ij} , $1 \leq i, j \leq n$ is a sum of commutators of the form $[u, v]$ ($u, v \in R$).

Remark 2.2. The proof of Theorem 2.1 yields that each c_{ij} , $1 \leq i, j \leq n$, $(i, j) \neq (1, 1)$ is a sum of commutators of the form $[\pm a', a'']$, where a' and a'' are products of certain entries of A .

Corollary 2.3. For the product $AA^* \in M_n(R)$ we have:

$$nAA^* = \operatorname{tr}(AA^*)I + C',$$

where $C' = [c'_{ij}]$ is an $n \times n$ matrix with $\operatorname{tr}(C') = 0$ and each c'_{ij} , $1 \leq i, j \leq n$ is a sum of commutators of the form $[u, v]$ ($u, v \in R$).

Proof. The claim easily follows from

$$\begin{aligned} C' &= nAA^* - \operatorname{tr}(AA^*)I \\ &= n(b_{11}I + C) - \operatorname{tr}(AA^*)I \\ &= (nb_{11} - \operatorname{tr}(AA^*))I + nC \\ &= ((b_{11} - b_{11}) + (b_{11} - b_{22}) + \cdots + (b_{11} - b_{nn}))I + nC \\ &= (-c_{22} - \cdots - c_{nn})I + nC. \end{aligned} \quad \square$$

Let $R[x]$ denote the ring of polynomials of the single commuting indeterminate x , with coefficients in R . The *right characteristic polynomial* of A is the right determinant of the $n \times n$ matrix $xI - A$ in $M_n(R[x])$:

$$p(x) = \operatorname{tr}((xI - A)(xI - A)^*) = \lambda_0 + \lambda_1x + \cdots + \lambda_{d-1}x^{d-1} + \lambda_d x^d \in R[x].$$

Proposition 2.4. If $p(x) = \lambda_0 + \lambda_1x + \cdots + \lambda_{d-1}x^{d-1} + \lambda_d x^d$ is the right characteristic polynomial of the $n \times n$ matrix $A \in M_n(R)$ then $d = n$ and $\lambda_d = n!$.

Proof. Take $(xI - A)^* = [h_{ij}(x)]$ and consider the trace of the product matrix $(xI - A)(xI - A)^*$:

$$\begin{aligned} p(x) &= \sum_{1 \leq i, j \leq n} (xI - A)_{ij} h_{ji}(x) \\ &= (x - a_{11})h_{11}(x) + \cdots + (x - a_{nn})h_{nn}(x) + \sum_{1 \leq i, j \leq n, i \neq j} (-a_{ij})h_{ji}(x). \end{aligned}$$

In view of the definition of the preadjoint, we can see that the degree of $h_{ij}(x)$ is $n - 2$ if $i \neq j$ and the leading monomial of $h_{ii}(x)$ is $(n - 1)!x^{n-1}$. Thus the leading monomial $n!x^n$ of $p(x)$ comes from

$$(x - a_{11})h_{11}(x) + \cdots + (x - a_{nn})h_{nn}(x). \quad \square$$

Proposition 2.5. *If the ring R is commutative, then we have*

$$p(x) = n! \det(xI - A)$$

for the right characteristic polynomial $p(x)$ of the $n \times n$ matrix $A \in M_n(R)$. Thus $p(x)$ is the $n!$ times scalar multiple of the ordinary characteristic polynomial of A .

Proof. Now we have

$$\begin{aligned} p(x) &= \operatorname{tr}((xI - A)(xI - A)^*) = \operatorname{tr}((xI - A)(n - 1)! \operatorname{adj}(xI - A)) \\ &= \operatorname{tr}((n - 1)! \det(xI - A)I) = n(n - 1)! \det(xI - A). \end{aligned}$$

We used the fact that now $(xI - A)^*$ can be expressed as

$$(xI - A)^* = (n - 1)! \operatorname{adj}(xI - A)$$

by the ordinary adjoint of $xI - A$ (see also in [9]). \square

3. The Cayley-Hamilton identity.

Theorem 3.1. *If $p(x) = \lambda_0 + \lambda_1 x + \cdots + \lambda_{n-1} x^{n-1} + n! x^n$ is the right characteristic polynomial of the $n \times n$ matrix $A \in M_n(R)$, then we can construct $n \times n$ matrices C_i , $0 \leq i \leq n$ with entries in $[R, R]$ such that*

$$(\lambda_0 I + C_0) + A(\lambda_1 I + C_1) + \cdots + A^{n-1}(\lambda_{n-1} I + C_{n-1}) + A^n(nI + C_n) = 0.$$

Proof. We use the natural isomorphism between the rings $M_n(R[x])$ and $M_n(R)[x]$. The application of Corollary 2.3 gives that

$$n(xI - A)(B_0 + B_1 x + \cdots + B_{n-1} x^{n-1}) = p(x)I + C'(x),$$

where

$$(xI - A)^* = [h_{ij}(x)] = B_0 + B_1 x + \cdots + B_{n-1} x^{n-1}$$

with $B_0, B_1, \dots, B_{n-1} \in M_n(R)$ (see the proof of Proposition 2.4) and $C'(x)$ is an $n \times n$ matrix with entries in $[R[x], R[x]]$ (and $\operatorname{tr}(C'(x)) = 0$). Since for

$f(x) = \sum_{\nu=1}^s u_\nu x^\nu$ and $g(x) = \sum_{\mu=1}^q v_\mu x^\mu$ in $R[x]$ the commutator

$$[f(x), g(x)] = f(x)g(x) - g(x)f(x) = \sum_{\nu, \mu} [u_\nu x^\nu, v_\mu x^\mu] = \sum_{\nu, \mu} [u_\nu, v_\mu] x^{\nu+\mu}$$

is a polynomial with coefficients in $[R, R]$, we can write that

$$C'(x) = C_0 + C_1x + \cdots + C_nx^n,$$

where C_0, C_1, \dots, C_n are $n \times n$ matrices with entries in $[R, R]$. The matching of the coefficients of the powers of x in the above matrix equation gives that

$$\begin{aligned} -nAB_0 &= \lambda_0I + C_0, \\ nB_0 - nAB_1 &= \lambda_1I + C_1, \\ &\vdots \\ nB_{n-2} - nAB_{n-1} &= \lambda_{n-1}I + C_{n-1}, \\ nB_{n-1} &= n!I + C_n. \end{aligned}$$

The left multiplication of $nB_{i-1} - nAB_i = \lambda_iI + C_i$ by A^i ($B_{-1} = B_n = 0$) gives the following sequence of matrix equations:

$$\begin{aligned} -nAB_0 &= \lambda_0I + C_0, \\ nAB_0 - nA^2B_1 &= A\lambda_1 + AC_1, \\ &\vdots \\ nA^{n-1}B_{n-2} - nA^nB_{n-1} &= A^{n-1}\lambda_{n-1} + A^{n-1}C_{n-1}, \\ nA^nB_{n-1} &= A^n n! + A^n C_n. \end{aligned}$$

Thus we obtain that

$$\begin{aligned} &(\lambda_0I + C_0) + A(\lambda_1I + C_1) + \cdots + A^{n-1}(\lambda_{n-1}I + C_{n-1}) + A^n(n!I + C_n) = \\ &= (-nAB_0) + (nAB_0 - nA^2B_1) + \cdots + (nA^{n-1}B_{n-2} - nA^nB_{n-1}) + (nA^nB_{n-1}) = 0. \square \end{aligned}$$

In view of the construction of the C_i 's in the above proof, it is reasonable to call

$$P(x) = n(xI - A)(xI - A)^* = p(x)I + C_0 + C_1x + \cdots + C_nx^n$$

the *generalized right characteristic polynomial* of $A \in M_n(R)$. Thus we have

$$np(x) = \text{tr}(P(x)).$$

Proposition 3.2. *If R is an algebra over a field K of characteristic zero and $T \in GL_n(K)$ then we have*

$$p_{TAT^{-1}}(x) = p_A(x) \text{ and } P_{TAT^{-1}}(x) = TP_A(x)T^{-1}$$

for the right characteristic polynomial $p_A(x) \in R[x]$ and the generalized right characteristic polynomial $P_A(x) \in M_n(R)[x]$ of $A \in M_n(R)$.

Proof. In [1] Domokos proved that $(TAT^{-1})^* = TA^*T^{-1}$, whence

$$\begin{aligned} P_{TAT^{-1}}(x) &= n(xI - TAT^{-1})(xI - TAT^{-1})^* \\ &= nT(xI - A)T^{-1}(T(xI - A)T^{-1})^* \\ &= nT(xI - A)T^{-1}T(xI - A)^*T^{-1} \\ &= Tn(xI - A)(xI - A)^*T^{-1} \\ &= TP_A(x)T^{-1} \end{aligned}$$

and

$$np_{TAT^{-1}}(x) = \text{tr}(TP_A(x)T^{-1}) = \text{tr}(P_A(x)) = np_A(x)$$

follows. Since $\frac{1}{n} \in K$, we conclude that $p_{TAT^{-1}}(x) = p_A(x)$. \square

REFERENCES

- [1] M. DOMOKOS. Cayley-Hamilton theorem for 2×2 matrices over the Grassmann algebra. *J. Pure Appl. Algebra* **133** (1998), 69–81.
- [2] A. R. KEMER. Varieties of \mathbb{Z}_2 -graded algebras. *Math. USSR Izv.* **25** (1985), 359–374.
- [3] A. R. KEMER. Ideals of Identities of Associative Algebras. Translations of Math. Monographs vol. **87**, 1991, AMS Providence, Rhode Island.
- [4] R. PARÉ, W. SCHELTER. Finite extensions are integral. *J. Algebra* **53** (1978), 477–479.
- [5] K. R. PEARSON. A lower bound for the degree of polynomials satisfied by matrices. *J. Aust. Math. Soc., Ser. A* **27** (1979), 430–436.
- [6] K. R. PEARSON. Degree 7 monic polynomials satisfied by a 3×3 matrix over a noncommutative ring. *Commun. Algebra* **10** (1982), 2043–2073.
- [7] K. R. PEARSON. The Robson cubics generate all polynomials satisfied by the general 2×2 matrix. *Commun. Algebra* **10** (1982), 2075–2084.

- [8] J. C. ROBSON. Polynomials satisfied by matrices. *J. Algebra* **55** (1978), 509–520.
- [9] J. SZIGETI. New determinants and the Cayley-Hamilton theorem for matrices over Lie nilpotent rings. *Proc. Amer. Math. Soc.* **125**, 8 (1997), 2245–2254.
- [10] J. SZIGETI. On the characteristic polynomial of supermatrices. *Israel J. Math.* **107** (1998), 229–235.

Institute of Mathematics

University of Miskolc

Miskolc, Hungary 3515

e-mail: jeno.szigeti@uni-miskolc.hu

Received July 15, 2006