

Provided for non-commercial research and educational use.
Not for reproduction, distribution or commercial use.

Serdica

Mathematical Journal

Сердика

Математическо списание

The attached copy is furnished for non-commercial research and education use only.
Authors are permitted to post this version of the article to their personal websites or institutional repositories and to share with other researchers in the form of electronic reprints.
Other uses, including reproduction and distribution, or selling or licensing copies, or posting to third party websites are prohibited.

For further information on
Serdica Mathematical Journal
which is the new series of
Serdica Bulgaricae Mathematicae Publicationes
visit the website of the journal <http://www.math.bas.bg/~serdica>
or contact: Editorial Office
Serdica Mathematical Journal
Institute of Mathematics and Informatics
Bulgarian Academy of Sciences
Telephone: (+359-2)9792818, FAX:(+359-2)971-36-49
e-mail: serdica@math.bas.bg

q -LEIBNIZ ALGEBRAS

A. S. Dzhumadil'daev

Communicated by V. Drensky

ABSTRACT. An algebra (A, \circ) is called Leibniz if $a \circ (b \circ c) = (a \circ b) \circ c - (a \circ c) \circ b$ for all $a, b, c \in A$. We study identities for the algebras $A^{(q)} = (A, \circ_q)$, where $a \circ_q b = a \circ b + qb \circ a$ is the q -commutator. Let $\text{char } K \neq 2, 3$. We show that the class of q -Leibniz algebras is defined by one identity of degree 3 if $q^2 \neq 1$, $q \neq -2$, by two identities of degree 3 if $q = -2$, and by the commutativity identity and one identity of degree 4 if $q = 1$. In the case of $q = -1$ we construct two identities of degree 5 that form a base of identities of degree 5 for -1 -Leibniz algebras. Any identity of degree < 5 for -1 -Leibniz algebras follows from the anti-commutativity identity.

1. Introduction. Denote by $A = (A, \circ)$ an algebra with vector space A over a field K of characteristic $\neq 2, 3$ and multiplication $(a, b) \mapsto a \circ b$. Let $(a, b, c) = a \circ (b \circ c) - (a \circ b) \circ c$ be the associator and $a \circ_q b = a \circ b + qb \circ a$ be the q -commutator, where $q \in K$. Denote by $A^{(q)} = (A, \circ_q)$ the algebra with the q -commutator. Notice that $a \circ_{-1} b = a \circ b - b \circ a$ is a commutator (Lie bracket,

2000 *Mathematics Subject Classification*: Primary 17A32, Secondary 17D25.

Key words: Leibniz algebras, Zinbiel algebras, Omni-Lie algebras, polynomial identities, q -commutators

usually denoted by $[a, b]$) and $a \circ_1 b = a \circ b + b \circ a$ is an anti-commutator (Jordan bracket, sometimes denoted by $\{a, b\}$).

Example. If A is an associative algebra, then $A^{(-1)} = (A, [,]) is a Lie algebra,$

$$[a, b] = -[b, a],$$

$$[[a, b], c] + [[b, c], a] + [[c, a], b] = 0,$$

and $A^{(+1)} = (A, \{ , \}) is a Jordan algebra,$

$$\{a, b\} = \{b, a\},$$

$$\{\{a, a\}, \{b, a\}\} = \{\{\{a, a\}, b\}, a\}.$$

Usually, q -commutators are studied in the frame of quantum groups. It seems that the study of q -identities has their own interest. We try to demonstrate it in the class of Leibniz algebras. We call an algebra A *Leibniz* (more exactly *right-Leibniz*) if for all $a, b, c \in A$

$$a \circ (b \circ c) = (a \circ b) \circ c - (a \circ c) \circ b.$$

Leibniz algebras were introduced in [2], [7]. In other words, Leibniz algebras are algebras with the identity $lei = 0$, where

$$lei = lei(t_1, t_2, t_3) = t_1(t_2t_3) - (t_1t_2)t_3 + (t_1t_3)t_2.$$

Example. Let (L, \star) be a Lie algebra with multiplication \star and let M be an L -module under the right action $(M, L) \rightarrow M, (m, a) \mapsto ma$. Make M a trivial left L -module: $am = 0, a \in L, m \in M$. Then the vector space $L \oplus M$ becomes a right-Leibniz algebra under the multiplication

$$(a + m) \circ (b + n) = a \star b + mb.$$

Indeed,

$$(a + m) \circ ((b + n) \circ (c + s)) = (a + m) \circ (b \star c + nc)$$

$$= a \star (b \star c) + m(b \star c) = (a \star b) \star c - (a \star c) \star b + (mb)c - (mc)b$$

$$= ((a + m) \circ (b + n)) \circ (c + s) - ((a + m) \circ (c + s)) \circ (b + n).$$

We call the so-obtained algebra $L + M$ (a semi-direct sum of Leibniz algebras) *standard Leibniz*.

Endow a standard Leibniz algebra $(L + M, \circ)$ with the commutator $[,]$.

Then

$$[a + m, b + n] = (a + m) \circ (b + n) - (b + n) \circ (a + m)$$

$$= (a \star b) + mb - (b \star a) - na = 2[a, b] + mb - na,$$

where $[a, b] = a \star b - b \star a$. The algebra $(L + M, [,])$ (more exactly, $L + M$ under multiplication $[a, b] + (mb - na)/2$) is called *Omni-Lie* [6], [9].

Given non-associative polynomials f_1, \dots, f_s , we let $\text{Var}(f_1, \dots, f_s)$ denote the variety of algebras defined by identities $f_1 = 0, \dots, f_s = 0$. Let \mathfrak{Lei} be the class of Leibniz algebras, i.e., the variety of algebras defined by the (right)-Leibniz identity $\text{lei} = 0$.

In this paper we construct identities for q -(right)-Leibniz algebras. In particular, we describe identities for Omni-Lie algebras.

We prove that the category of q -Leibniz algebras is equivalent to the category of Leibniz algebras if $q^2 \neq 1, q \neq -2$. This means that, for $q \neq \pm 1, -2$, every algebra with identity $\text{lei}^{(q)} = 0$ can be obtained as $A^{(q)}$ from some Leibniz algebra A and, conversely, if B is an algebra with identity $\text{lei}^{(q)} = 0$, then $B^{(-q)}$ is right-Leibniz. In the case of $q = -2$ we should add to the identity $\text{lei}^{(q)} = 0$ the identity $\text{lei}_1^{(q)} = 0$ in order to obtain equivalent categories.

Theorem 1.1. *Let $q \neq -1, 1, -2$. The class of q -Leibniz algebras $\mathfrak{Lei}^{(q)}$ satisfies the identity $\text{lei}^{(q)} = 0$, where*

$$\text{lei}^{(q)} = \text{lei}^{(q)}(t_1, t_2, t_3)$$

$$= (q^2 - 1)(t_1(t_2t_3) - t_2(t_1t_3)) + (q^2 + q - 1)(t_2t_1)t_3 + (t_2t_3)t_1 - t_1(t_3t_2) - qt_3(t_1t_2).$$

The varieties \mathfrak{Lei} , $\mathfrak{Lei}^{(q)}$ and $\text{Var}(\text{lei}^{(q)})$ are equivalent.

In particular, $\text{Var}(\text{lei}^{(q)})$ has no special identity for $\mathfrak{Lei}^{(q)}$ if $q \neq -2, q^2 \neq 1$. The identity $\text{lei}_1^{(q)} = 0$ is a consequence of the identity $\text{lei}^{(q)} = 0$ if $q \neq -2, q^2 \neq 1$.

Theorem 1.2. *Let $q = -2$. The class of q -Leibniz algebras $\mathfrak{Lei}^{(-2)}$ satisfies the identities $\text{lei}^{(-2)} = 0$ and $\text{lei}_1^{(-2)} = 0$, where $\text{lei}^{(q)}$ is given above and*

$$\text{lei}_1^{(q)} = \text{lei}_1^{(q)}(t_1, t_2, t_3) = -t_1(t_2t_3 + t_3t_2) + q(t_2t_3 + t_3t_2)t_1.$$

The varieties \mathfrak{Lei} , $\mathfrak{Lei}^{(-2)}$ and $\text{Var}(\text{lei}^{(-2)}, \text{lei}_1^{(-2)})$ are equivalent.

So the identity $\text{lei}_1^{(-2)} = 0$ is a special identity for $\text{Var}(\text{lei}^{(-2)})$ which does not follow from the identity $\text{lei}^{(-2)} = 0$. All other special identities for $\mathfrak{Lei}^{(-2)}$ follow from $\text{lei}^{(-2)} = 0$.

Let acom , com and ljac be non-commutative non-associative polynomials defined by

$$\text{acom} = t_1t_2 + t_2t_1,$$

$$\text{com} = t_1t_2 - t_2t_1,$$

$$\text{ljac} = (t_1t_2)t_3 + (t_2t_3)t_1 + (t_3t_1)t_2.$$

Define non-commutative non-associative polynomials $leilie_1$, $leilie_2$ of degree five by

$$leilie_1(t_1, t_2, t_3, t_4, t_5) = 2\text{ljac}(\text{ljac}(t_1, t_2, t_3), t_4, t_5) - [\text{ljac}(t_1, t_2, t_3), [t_4, t_5]],$$

$$\begin{aligned} leilie_2(t_1, t_2, t_3, t_4, t_5) = & -\frac{1}{2} \sum_{\sigma \in \text{Sym}(2,3,4,5)} \text{sign } \sigma (-4((t_{\sigma(2)}t_{\sigma(3)})t_{\sigma(4)})t_{\sigma(5)})t_1 \\ & + 2(((t_{\sigma(2)}t_{\sigma(3)})t_1)t_{\sigma(4)})t_{\sigma(5)} + 2(((t_{\sigma(2)}t_{\sigma(3)})t_{\sigma(4)})t_1)t_{\sigma(5)} \\ & + ((t_1t_{\sigma(2)})t_{\sigma(3)})(t_{\sigma(4)}t_{\sigma(5)}) + ((t_1t_{\sigma(2)})(t_{\sigma(4)}t_{\sigma(5)}))t_{\sigma(3)}. \end{aligned}$$

For a non-commutative non-associative polynomial $f(t_1, \dots, t_k)$, denote by $\text{Alt}(f)$ its skew-symmetrization

$$\text{Alt } f(t_1, \dots, t_k) = \sum_{\sigma \in \text{Sym}_k} \text{sign } \sigma f(t_{\sigma(1)}, \dots, t_{\sigma(k)}).$$

Let

$$leilie(t_1, t_2, t_3, t_4, t_5) = \text{Alt}(4((t_1t_2)t_3)t_4)t_5 - ((t_1t_2)t_3)(t_4t_5).$$

Theorem 1.3. *Let $q = -1$. Let A be a right-Leibniz algebra. Then $A^{(-1)}$ satisfies the identities $\text{acom} = 0$, $leilie_1 = 0$ and $leilie_2 = 0$. Any multilinear identity of $\mathfrak{Lci}^{(-1)}$ of degree no more than 4 follows from the anti-commutativity identity. Any multilinear identity of $\mathfrak{Lci}^{(-1)}$ of degree 5 follows from the identities $\text{acom} = 0$, $leilie_1 = 0$ and $leilie_2 = 0$.*

Corollary 1.4. *Let A be a right-Leibniz algebra. Then $A^{(-1)}$ satisfies the identity $leilie = 0$.*

Corollary 1.5. *Every Omni-Lie algebra satisfies the polynomial identities $\text{acom} = 0$, $leilie_1 = 0$, $leilie_2 = 0$ and $leilie = 0$. The identities $\text{acom} = 0$, $leilie_1 = 0$ and $leilie_2 = 0$ form a base of the identities in the space of multilinear identities of degree no more than 5 for the class of Omni-Lie algebras.*

Note that the polynomials $leilie_1$, $leilie_2$ and $leilie$ have 9, 60 and 90 terms, respectively.

Let

$$leijor(t_1, t_2, t_3, t_4) = (t_1t_2)(t_3t_4).$$

Theorem 1.6. *Let $q = 1$. Let A be a right-Leibniz algebra. Then $A^{(1)}$ satisfies the identities $\text{com} = 0$ and $leijor = 0$. Every multilinear identity which*

is true for any Leibniz-Jordan algebra follows from the identities $\text{com} = 0$ and $\text{leijor} = 0$.

In other words, there are no special identities for the class of Leibniz-Jordan algebras.

The Leibniz operad has a dual operad defined by the identity

$$a(bc + cb) = (ab)c + (ac)b.$$

Such algebras are called Zinbiel [7], [8]. Identities for *q*-Zinbiel algebras are described in [3], [4].

2. Non-commutative non-associative polynomials. Let $K\{t_1, t_2, \dots\}$ be the algebra of non-commutative non-associative polynomials in the variables t_1, t_2, \dots (the free magma algebra). For a polynomial $f = f(t_1, \dots, t_k) \in K\{t_1, t_2, \dots\}$, we say that $f = 0$ is an *identity* for the algebra (A, \circ) if $f(a_1, \dots, a_k) = 0$ for all $a_1, \dots, a_k \in A$.

Recall that there exist $\frac{1}{k} \binom{2(k-1)}{k-1}$ types of bracketing for the string $t_1 \cdots t_k$. For example, there are 5 types of bracketing for 4 elements:

$$((t_1 t_2) t_3) t_4, (t_1 t_2)(t_3 t_4), t_1(t_2(t_3 t_4)), t_1((t_2 t_3) t_4), (t_1(t_2 t_3)) t_4.$$

Order the types of bracketing somehow. If σ is a type of bracketing, denote by $\sigma(t_{i_1}, \dots, t_{i_k})$ the string $t_{i_1} \cdots t_{i_k}$ with bracketing type σ . For example, if $k = 4$ and σ is the bracketing type $(t_1(t_2 t_3)) t_4$ then $\sigma(t_1, t_2, t_1, t_3) = (t_1(t_2 t_1)) t_3$.

Let α be some bracketing type of t_1, \dots, t_n . We say that a monomial of the form $\alpha(t_{i_1}, \dots, t_{i_n})$ has *multidegree* (r_1, \dots, r_k) if $\{i_1, \dots, i_n\} \subseteq \{1, \dots, k\}$ and $r_m = |\{s : i_s = m, s = 1, \dots, n\}|$ is the number of indices i_s equal to m for any $m = 1, \dots, k$. Call $f = f(x_1, \dots, x_k)$ *homogeneous of degree* (r_1, \dots, r_k) if f is a linear combination of monomials of multidegree (r_1, \dots, r_k) . Say that a homogeneous polynomial f has *degree* l if $r_1 + \dots + r_k = l$.

A homogeneous polynomial $f = f(t_1, \dots, t_k)$ of multidegree $(1, \dots, 1)$ is called *multilinear*. Notice that the degree of a multilinear polynomial $f \in K\{t_1, \dots, t_k\}$ is equal to the number of variables k . In other words a polynomial f is multilinear if f is a linear combination of monomials of the form $\alpha(t_{i_1}, \dots, t_{i_k})$, where $\begin{pmatrix} 1 & \cdots & k \\ i_1 & \cdots & i_k \end{pmatrix} \in \text{Sym}_k$ is a permutation of the set $\{1, \dots, k\}$ and α is a bracketing.

Given polynomials $f_1, \dots, f_s, g \in K\{t_1, \dots, t_k\}$, we say that the identity $g = 0$ follows from the identities $f_1 = 0, \dots, f_s = 0$, and write $\{f_1 = 0, \dots, f_s =$

$0\} \Rightarrow g = 0$, if $g = 0$ is an identity for any algebra in the variety defined by the identities $f_1 = 0, \dots, f_s = 0$.

Let \mathfrak{L} be a variety of algebras and let $\mathfrak{L}^{(q)}$ be the class of algebras $A^{(q)}$ such that $A \in \mathfrak{L}$. Suppose that $(A, \circ_q) \in \mathfrak{L}^{(q)}$ has identities $f_1 = 0, \dots, f_s = 0$. We say that these identities are $\mathfrak{L}^{(q)}$ -minimal if

- for any $r = 1, \dots, s$, the identity $f_r = 0$ does not follow from the identities $f_1 = 0, \dots, f_{r-1} = 0, f_{r+1} = 0, \dots, f_s = 0$;
- if $\{f_1 = 0, \dots, f_{r-1} = 0, g = 0, f_{r+1} = 0, \dots, f_s = 0\} \Rightarrow f_r = 0$ and $g = 0$ is an identity for $\mathfrak{L}^{(q)}$ then $\{f_1 = 0, \dots, f_{r-1} = 0, f_r = 0, f_{r+1} = 0, \dots, f_s = 0\} \Rightarrow g = 0$.

Let $(f, g) \rightarrow f \cdot g = fg$ be the multiplication of the algebra $K\{t_1, t_2, \dots\}$. Let us endow the algebra with the multiplication $(f, g) \mapsto f \cdot_q g$ given by $f \cdot_q g = f \cdot g + qg \cdot f$. For example,

$$\begin{aligned} (t_1 + 3t_1t_2) \cdot ((t_2t_3)t_1) &= t_1((t_2t_3)t_1) + 3(t_1t_2)((t_2t_3)t_1), \\ (t_1 + 3t_1t_2) \cdot_q ((t_2t_3)t_1) &= t_1((t_2t_3)t_1) + 3(t_1t_2)((t_2t_3)t_1) \\ &\quad + q((t_2t_3)t_1)t_1 + 3q((t_2t_3)t_1)(t_1t_2). \end{aligned}$$

Let

$$\tau_q : K\{t_1, t_2, \dots\} \rightarrow K\{t_1, t_2, \dots\}$$

be a linear map defined by

$$\begin{aligned} \tau_q(t_i) &= t_i, \\ \tau_q(f \cdot g) &= \tau_q(f) \cdot \tau_q(g) + q \tau(g) \cdot \tau_q(f), \end{aligned}$$

for any $f, g \in K\{t_1, t_2, \dots\}$. Then

$$\tau_q : (K\{t_1, t_2, \dots\}, \cdot) \rightarrow (K\{t_1, t_2, \dots\}, \cdot_q)$$

is the homomorphism

$$\tau_q(f \cdot g) = \tau_q(f) \cdot_q \tau_q(g).$$

Given a bracketing type σ , we set

$$\sigma_q = \tau_q \sigma.$$

In other words, $\sigma_q(t_1, \dots, t_k)$ is the polynomial obtained from $\sigma(t_1, \dots, t_k)$ by the multiplication \circ_q . For example, if σ is the bracketing type $(t_1t_2)t_3$, then

$$\sigma_q(t_3, t_1, t_2) = (t_3t_1)t_2 + q((t_1t_3)t_2 + t_2(t_3t_1)) + q^2t_2(t_1t_3).$$

Lemma 2.1. For any bracketing type σ

$$\sigma_{-q} \sigma_q(t_{i_1}, \dots, t_{i_k}) = (1 - q^2)^{k-1} \sigma_0(t_{i_1}, \dots, t_{i_k}).$$

Proof. We use induction on *k*. For *k* = 2 the statement is true:

$$\sigma_q(t_{i_1}, t_{i_2}) = t_{i_1}t_{i_2} + q t_{i_2}t_{i_1},$$

and

$$\begin{aligned} \sigma_{-q}\sigma_q(t_{i_1}, t_{i_2}) &= t_{i_1}t_{i_2} - q t_{i_2}t_{i_1} + q t_{i_2} \cdot t_{i_1} - q^2 t_{i_1}t_{i_2} \\ &= (1 - q^2)t_{i_1}t_{i_2} = (1 - q^2)\sigma_0(t_{i_1}, t_{i_2}). \end{aligned}$$

Suppose that our statement is true for *k* - 1. Let

$$\sigma(t_{i_1}, \dots, t_{i_k}) = \sigma'(t_{i_1}, \dots, t_{i_{k'}})\sigma''(t_{i_{k'+1}}, \dots, t_{i_k})$$

for some $1 \leq k' \leq k$ and for some bracketings σ', σ'' . Then

$$\begin{aligned} \sigma_q(t_{i_1}, \dots, t_{i_k}) &= \sigma'_q(t_{i_1}, \dots, t_{i_{k'}})\sigma''_q(t_{i_{k'+1}}, \dots, t_{i_k}) \\ &\quad + q \sigma''_q(t_{i_{k'+1}}, \dots, t_{i_k})\sigma'_q(t_{i_1}, \dots, t_{i_{k'}}) \end{aligned}$$

and

$$\begin{aligned} \sigma_{-q}\sigma_q(t_{i_1}, \dots, t_{i_k}) &= \sigma'_{-q}\sigma'_q(t_{i_1}, \dots, t_{i_{k'}})\sigma''_{-q}\sigma''_q(t_{i_{k'+1}}, \dots, t_{i_k}) \\ &\quad - q \sigma''_{-q}\sigma''_q(t_{i_{k'+1}}, \dots, t_{i_k})\sigma'_{-q}\sigma'_q(t_{i_1}, \dots, t_{i_{k'}}) \\ &\quad + q \sigma''_{-q}\sigma''_q(t_{i_{k'+1}}, \dots, t_{i_k})\sigma'_{-q}\sigma'_q(t_{i_1}, \dots, t_{i_{k'}}) \\ &\quad - q^2 \sigma'_{-q}\sigma'_q(t_{i_1}, \dots, t_{i_{k'}})\sigma''_{-q}\sigma''_q(t_{i_{k'+1}}, \dots, t_{i_k}) \\ &= \sigma'_{-q}\sigma'_q(t_{i_1}, \dots, t_{i_{k'}})\sigma''_{-q}\sigma''_q(t_{i_{k'+1}}, \dots, t_{i_k}) \\ &\quad - q^2 \sigma'_{-q}\sigma'_q(t_{i_1}, \dots, t_{i_{k'}})\sigma''_{-q}\sigma''_q(t_{i_{k'+1}}, \dots, t_{i_k}). \end{aligned}$$

By the induction hypothesis

$$\begin{aligned} \sigma'_{-q}\sigma'_q(t_{i_1}, \dots, t_{i_{k'}}) &= (1 - q^2)^{k'-1}\sigma'_0(t_{i_1}, \dots, t_{i_{k'}}), \\ \sigma''_{-q}\sigma''_q(t_{i_{k'+1}}, \dots, t_{i_k}) &= (1 - q^2)^{k-k'-1}\sigma''_0(t_{i_{k'+1}}, \dots, t_{i_k}). \end{aligned}$$

Therefore,

$$\begin{aligned} \sigma'_{-q}\sigma'_q(t_{i_1}, \dots, t_{i_{k'}})\sigma''_{-q}\sigma''_q(t_{i_{k'+1}}, \dots, t_{i_k}) &= (1 - q^2)^{k-2}\sigma_0(t_{i_1}, \dots, t_{i_k}), \\ -q^2 \sigma'_{-q}\sigma'_q(t_{i_1}, \dots, t_{i_{k'}})\sigma''_{-q}\sigma''_q(t_{i_{k'+1}}, \dots, t_{i_k}) &= -q^2(1 - q^2)^{k-2}\sigma_0(t_{i_1}, \dots, t_{i_k}) \end{aligned}$$

and

$$\begin{aligned} \sigma_{-q}\sigma_q(t_{i_1}, \dots, t_{i_k}) &= \sigma'_{-q}\sigma'_q(t_{i_1}, \dots, t_{i_{k'}})\sigma''_{-q}\sigma''_q(t_{i_{k'+1}}, \dots, t_{i_k}) \\ &\quad - q^2 \sigma'_{-q}\sigma'_q(t_{i_1}, \dots, t_{i_{k'}})\sigma''_{-q}\sigma''_q(t_{i_{k'+1}}, \dots, t_{i_k}) \\ &= (1 - q^2)^{k-1}\sigma_0(t_{i_1}, \dots, t_{i_k}). \end{aligned}$$

From Lemma 2.1 we infer the following

Theorem 2.2. $(q^2 \neq 1)$ Let f_1, \dots, f_s be homogeneous polynomials of degree k . Then the class of q -algebras $\text{Var}(f_1, \dots, f_s)^{(q)}$ forms a variety defined by the system of polynomial identities $\sigma_{-q}f_1 = 0, \dots, \sigma_{-q}f_s = 0$. This variety is equivalent to $\text{Var}(f_1, \dots, f_s)$ and the equivalence can be given by $A = (A, \star) \mapsto A^{(-q)} = (A, \star_{-q})$.

The equivalence of varieties means the following. There exist functors

$$F : \text{Var}(f_1, \dots, f_s) \rightarrow \text{Var}(\sigma_{-q}f_1, \dots, \sigma_{-q}f_s), \quad (A, \circ) \rightarrow (A, \circ_q),$$

$$G : \text{Var}(\sigma_{-q}f_1, \dots, \sigma_{-q}f_s) \rightarrow \text{Var}(f_1, \dots, f_s), \quad (A, \star) \rightarrow (A, \star'_q)$$

such that

$$GF(A, \circ) = (A, \circ), \quad FG(A, \star) = (A, \star).$$

Here

$$a \star'_q b = \frac{1}{(1 - q^2)^{k-1}} a \star_q b.$$

Recall that all polynomials f_1, \dots, f_s are supposed homogeneous. Notice that, for any $(A, \circ), (B, \cdot) \in \text{Var}(f_1, \dots, f_s)$ and a morphism between them, i.e., a homomorphism $\psi : (A, \circ) \rightarrow (B, \cdot)$, there corresponds a morphism of algebras $\psi : F(A, \circ) \rightarrow F(B, \cdot)$ in the category $\text{Var}(\sigma_{-q}f_1, \dots, \sigma_{-q}f_s)$, i.e., a homomorphism $\psi : (A, \circ_q) \rightarrow (B, \cdot_q)$. Indeed,

$$\begin{aligned} \psi(a_1 \circ_q a_2) &= \psi(a_1 \circ a_2 + q a_2 \circ a_1) \\ &= \psi(a_1 \circ a_2) + q \psi(a_2 \circ a_1) \\ &= \psi(a_1) \cdot \psi(a_2) + q \psi(a_2) \cdot \psi(a_1) \\ &= \psi(a_1) \cdot_q \psi(a_2). \end{aligned}$$

If I is an ideal of (A, \circ) then I is an ideal of (A, \circ_q) . Therefore, simplicity, nilpotency and solvability properties of algebras in the category $\text{Var}(f_1, \dots, f_s)$ remain the same for the corresponding algebras in $\text{Var}(\sigma_{-q}f_1, \dots, \sigma_{-q}f_s)$. If (A, \circ) is free in the variety $\text{Var}(f_1, \dots, f_s)$, then (A, \circ_q) is free in the variety $\text{Var}(\sigma_{-q}f_1, \dots, \sigma_{-q}f_s)$. We pay attention to the fact that the categories $\text{Var}(f_1, \dots, f_s)$ and $\text{Var}(\sigma_{-q}f_1, \dots, \sigma_{-q}f_s)$ are equivalent only in the case of $q^2 \neq 1$.

Let g_1, \dots, g_s, h be non-commutative non-associative polynomials. Suppose that, for a class \mathfrak{L} of algebras, the corresponding class $\mathfrak{L}^{(q)}$ of q -algebras satisfies the identities $g_1 = 0, \dots, g_s = 0$ and $h = 0$. In this case we say that $h = 0$ is a *special* identity or an *s*-identity for $\text{Var}(g_1, \dots, g_s)$.

We give another application of Lemma 2.1.

Theorem 2.3. *If $q \neq \pm 1$, then the map*

$$\tau_q : (K\{t_1, t_2, \dots\}, \cdot) \rightarrow (K\{t_1, t_2, \dots\}, \cdot_q)$$

is an isomorphism.

Let \mathfrak{L} be some class of algebras. For a polynomial $f \in K\{t_1, t_2, \dots\}$, we say that $f = 0$ is an identity for \mathfrak{L} if every algebra $A \in \mathfrak{L}$ satisfies the identity $f = 0$. Recall that the class of all algebras satisfying given polynomial identities forms a variety.

Recall that \mathfrak{Lei} is the class of Leibniz algebras and $\mathfrak{Lei}^{(q)}$ is the class of *q*-Leibniz algebras, i.e., algebras of the form $A^{(q)} = (A, \circ_q)$, where $A \in \mathfrak{Lei}$.

Define non-commutative polynomials *rjac* (*right-Jacobian*), *lalia* (*left-anti-Lie-admissible*), *ralia* (*right-Anti-Lie-admissible*), *lia* (*Lie-admissible*), s_k^l (*standard left-skew-symmetric*), s_k^r (*standard right-skew-symmetric*) and $s_k^{[r]}$ (*s_k-Lie-admissible*) by

$$\begin{aligned} \text{rjac}(t_1, t_2, t_3) &= t_1(t_2t_3) + t_2(t_3t_1) + t_3(t_1t_2), \\ \text{lalia}(t_1, t_2, t_3) &= [t_1, t_2]t_3 + [t_2, t_3]t_1 + [t_3, t_1]t_2, \\ \text{ralia}(t_1, t_2, t_3) &= t_1[t_2, t_3] + t_2[t_3, t_1] + t_3[t_1, t_2], \\ \text{lia}(t_1, t_2, t_3) &= [[t_1, t_2], t_3] + [[t_2, t_3], t_1] + [[t_3, t_1], t_2], \\ \text{alia}^{(q)} &= \text{lalia} + q \cdot \text{ralia}, \quad q \in K. \end{aligned}$$

Recall that for a non-commutative non-associative polynomial $f(t_1, \dots, t_k)$, we denote by $\text{Alt}(f)$ its skew-symmetrization

$$\text{Alt } f(t_1, \dots, t_k) = \sum_{\sigma \in \text{Sym}_k} \text{sign } \sigma f(t_{\sigma(1)}, \dots, t_{\sigma(k)}).$$

Let

$$\begin{aligned} s_k^r(t_1, \dots, t_k) &= \text{Alt}(t_1(t_2(\dots(t_{k-1}t_k))))), \\ s_k^l(t_1, \dots, t_k) &= \text{Alt}((\dots(t_1t_2)\dots t_{k-1})t_k), \\ s_k^{[r]}(t_1, \dots, t_k) &= \text{Alt}([t_1, [t_2, \dots, [t_{l-1}, t_k]]]). \end{aligned}$$

Notice that

$$\text{com} = s_2, \quad \text{lalia} = s_3^l, \quad \text{ralia} = s_3^r, \quad \text{lia} = s_3^l - s_3^r = \text{lalia} - \text{ralia}.$$

If algebras are anti-commutative, i.e., satisfy the identity $\text{acom} = 0$, then

$$\text{ljac} = -\text{rjac},$$

$$\text{lia} = 4 \text{ljac}.$$

3. Right-center and Lie elements. Let $F = F(V)$ be a free right-Leibniz algebra generated by a space V . Let $(F^{\text{lie}}, [,])$ be the subspace of F generated by V under the commutator $[,]$. We say that $a \in F$ is a *Lie-element* if $a \in (F^{\text{lie}}, [,])$. Homomorphic images of Lie elements of any Leibniz algebras are called Lie elements as well.

Let (A, \circ) be a right-Leibniz algebra. An element $z \in A$ is called *right-central* if

$$a \circ z = 0$$

for all $a \in A$. Let A^{rann} be the set of right-central elements of A . It was noticed in [7] that A^{rann} is an ideal with trivial left action, $a \circ z = 0, z \in A^{\text{rann}}, a \in A$, such that

$$\{a, b\} = a \circ b + b \circ a \in A^{\text{rann}}$$

for all $a, b \in A$. We construct new right-central elements.

Observe that

$$(1) \quad s_{k+1}^l(a_1, \dots, a_{k+1}) = \sum_{i=1}^{k+1} (-1)^i s_k^l(a_1, \dots, \hat{a}_i, \dots, a_{k+1}) \circ a_i.$$

Lemma 3.1. *Let (A, \circ) be a right-Leibniz algebra. Then A^{rann} is an ideal such that*

$$a \circ z = 0.$$

For any $q \in K$,

$$(2) \quad (a \circ_q b) \circ c = (a \circ c) \circ_q b + a \circ_q (b \circ c).$$

In particular,

$$\{a, b\} \circ c = \{a \circ c, b\} + \{a, b \circ c\}.$$

For any $k \geq 3$

$$s_k^l(a_1, \dots, a_k), \quad s_k^r(a_1, \dots, a_k) \in A^{\text{rann}}.$$

Moreover,

$$s_k^l(a_1, \dots, a_k) = s_k^{[r]}(a_1, \dots, a_k)$$

are Lie elements,

$$s_k^r(a_1, \dots, a_k) = 0, k \geq 4,$$

and

$$s_3^r(a, b, c) = 2s_3^l(a, b, c).$$

In other words, any right-Leibniz algebra *A* is $-1/2$ -Alia, i.e.,

$$\text{alia}^{(-1/2)}(a, b, c) = 0$$

for all $a, b, c \in A$.

Proof. We have

$$\begin{aligned} (a \circ_q b) \circ c &= (a \circ b + qb \circ a) \circ c \\ &= a \circ (b \circ c) + (a \circ c) \circ b + qb \circ (a \circ c) + q(b \circ c) \circ a \\ &= (a \circ c) \circ_q b + a \circ_q (b \circ c). \end{aligned}$$

So, (2) is established. Thus, in the case $q = 0$ we obtain the right-Leibniz identity

$$(a \circ b) \circ c = (a \circ c) \circ b + a \circ (b \circ c).$$

Let $k = 3$. Notice that

$$s_3^r(a, b, c) = \text{ralia}(a, b, c).$$

We have

$$\begin{aligned} \text{ralia}(a, b, c) &= a \circ [b, c] + b \circ [c, a] + c \circ [a, b] \\ &= 2(a \circ (b \circ c) + b \circ (c \circ a) + c \circ (a \circ b)) \\ &\quad - a \circ \{b, c\} - b \circ \{c, a\} - c \circ \{a, b\} \\ &= 2 \text{rjac}(a, b, c). \end{aligned}$$

By the right-Leibniz identity

$$\text{ljac}(a, b, c) = [a, b] \circ c + [b, c] \circ a + [c, a] \circ b = \text{lalia}(a, b, c),$$

and

$$\begin{aligned} \text{lia}(a, b, c) &= \text{lalia}(a, b, c) - \text{ralia}(a, b, c) = \text{rjac}(a, b, c) - 2 \text{rjac}(a, b, c) \\ &= - \text{rjac}(a, b, c). \end{aligned}$$

So, $s_3^r(a, b, c) = 2\text{rjac}(a, b, c) = -2\text{lia}(a, b, c)$ is a Lie-element.

By the right-Leibniz identity

$$\begin{aligned} u \circ \text{rjac}(a, b, c) &= ((u \circ a) \circ (b \circ c) - ((u \circ (b \circ c)) \circ a + ((u \circ b) \circ (c \circ a) \\ &\quad - ((u \circ (c \circ a)) \circ b + ((u \circ c) \circ (a \circ b) - ((u \circ (a \circ b)) \circ c \\ &= ((u \circ a) \circ b) \circ c - ((u \circ a) \circ c) \circ b - ((u \circ b) \circ c) \circ a \\ &\quad + ((u \circ c) \circ b) \circ a + ((u \circ b) \circ c) \circ a - ((u \circ b) \circ a) \circ c \\ &\quad - ((u \circ c) \circ a) \circ b + ((u \circ a) \circ c) \circ b + ((u \circ c) \circ a) \circ b \\ &\quad - ((u \circ c) \circ b) \circ a - ((u \circ a) \circ b) \circ c + ((u \circ b) \circ a) \circ c = 0. \end{aligned}$$

So, the element $s_3^l(a, b, c)$ is right-central.

Suppose that $s_k^l(a_1, \dots, a_k) = s^{[r]}(a_1, \dots, a_k)$ is a Lie element and is right-central. Prove that $s_{k+1}^l(a_1, \dots, a_{k+1})$ is also a Lie element which is right-central. Since $s_k^l(a_1, \dots, \hat{a}_i, \dots, a_{k+1}) \in A^{\text{rann}}$ for every $i = 1, \dots, k+1$ and since A^{rann} is an ideal, we have

$$s_{k+1}^l(a_1, \dots, a_{k+1}) = \sum_{i=1}^{k+1} (-1)^{i+k+1} s_k^l(a_1, \dots, \hat{a}_i, \dots, a_{k+1}) \circ a_i \in A^{\text{rann}}.$$

Further,

$$s_{k+1}^{[r]}(a_1, \dots, a_{k+1}) = \sum_{i=1}^{k+1} (-1)^{i+1} [a_i, s_k^{[r]}(a_1, \dots, \hat{a}_i, \dots, a_{k+1})]$$

(by the induction hypothesis)

$$= \sum_{i=1}^{k+1} (-1)^{i+1} [a_i, s_k^l(a_1, \dots, \hat{a}_i, \dots, a_{k+1})]$$

(since $s_k^l(a_1, \dots, \hat{a}_i, \dots, a_{k+1}) \in A^{\text{rann}}$)

$$= \sum_{i=1}^{k+1} (-1)^i s_k^l(a_1, \dots, \hat{a}_i, \dots, a_{k+1}) \circ a_i = s_{k+1}^l(a_1, \dots, a_{k+1}).$$

4. *q*-commutators of Leibniz algebras in case $q^2 \neq 1$.

Lemma 4.1. *For any Leibniz algebra A its q -algebra $A^{(q)}$ satisfies the identities $\text{lei}^{(q)} = 0$ and $\text{lei}_1^{(q)} = 0$.*

Proof. We have

$$\begin{aligned} \text{lei}^{(q)}(a, b, c) &= (q^2 - 1) a \circ_q (b \circ_q c) - a \circ_q (c \circ_q b) - (q^2 - 1) b \circ_q (a \circ_q c) \\ &\quad - qc \circ_q (a \circ_q b) + (q^2 + q - 1) (b \circ_q a) \circ_q c + (b \circ_q c) \circ_q a \\ &= (q^2 - 1)(a \circ (b \circ c) + (1 + q)a \circ (c \circ b) - b \circ (a \circ c) - qb \circ (c \circ a) \\ &\quad + (q + q^2)c \circ (a \circ b) + qc \circ (b \circ a) + q(a \circ b) \circ c - q(a \circ c) \circ b \\ &\quad + (1 - q)(b \circ a) \circ c + (q - 1)(b \circ c) \circ a - q^2(c \circ a) \circ b \\ &\quad + q^2(c \circ b) \circ a) \\ &= (q^2 - 1)(a \circ (b \circ c) + (1 + q)a \circ (c \circ b) - b \circ (a \circ c) + qb \circ (a \circ c) \\ &\quad + q^2c \circ (a \circ b) + q(a \circ b) \circ c - q(a \circ c) \circ b + (1 - q)(b \circ a) \circ c \\ &\quad + (q - 1)(b \circ c) \circ a - q^2(c \circ a) \circ b + q^2(c \circ b) \circ a) \\ &= (q^2 - 1)(qa \circ (c \circ b) + (q - 1)b \circ (a \circ c) + q^2c \circ (a \circ b) \\ &\quad + q((a \circ b) \circ c - (a \circ c) \circ b) + (1 - q)((b \circ a) \circ c - (b \circ c) \circ a) \\ &\quad - q^2((c \circ a) \circ b - (c \circ b) \circ a)) \\ &= (q^2 - 1)(q(a \circ (c \circ b) + (a \circ b) \circ c - (a \circ c) \circ b) \\ &\quad + (1 - q)(-b \circ (a \circ c) + (b \circ a) \circ c - (b \circ c) \circ a) \\ &\quad - q^2(-c \circ (a \circ b) + (c \circ a) \circ b - (c \circ b) \circ a)) \end{aligned}$$

(by the right-Leibniz identity)

$$= 0.$$

Similarly,

$$\begin{aligned}
\text{lei}_1^{(q)}(a, b, c) &= -a \circ_q (b \circ_q c) - a \circ_q (c \circ_q b) + q(b \circ_q c) \circ_q a + q(c \circ_q b) \circ_q a \\
&= -a \circ (b \circ c) - qa \circ (c \circ b) - q(b \circ c) \circ a - q^2(c \circ b) \circ a \\
&\quad - a \circ (c \circ b) - qa \circ (b \circ c) - q(c \circ b) \circ a - q^2(b \circ c) \circ a \\
&\quad + q(b \circ c) \circ a + q^2(c \circ b) \circ a + q^2a \circ (b \circ c) + q^3a \circ (c \circ b) \\
&\quad + q(c \circ b) \circ a + q^2(b \circ c) \circ a + q^2a \circ (c \circ b) + q^3a \circ (b \circ c) \\
&= -(a \circ b) \circ c + (a \circ c) \circ b - q(a \circ c) \circ b + q(a \circ b) \circ c \\
&\quad - q(b \circ c) \circ a - q^2(c \circ b) \circ a \\
&\quad - (a \circ c) \circ b + (a \circ b) \circ c - q(a \circ b) \circ c + q(a \circ c) \circ b \\
&\quad - q(c \circ b) \circ a - q^2(b \circ c) \circ a + q(b \circ c) \circ a + q^2(c \circ b) \circ a \\
&\quad + q^2(a \circ b) \circ c - q^2(a \circ c) \circ b + q^3(a \circ c) \circ b - q^3(a \circ b) \circ c \\
&\quad + q(c \circ b) \circ a + q^2(b \circ c) \circ a \\
&\quad + q^2(a \circ c) \circ b - q^2(a \circ b) \circ c + q^3(a \circ b) \circ c - q^3(a \circ c) \circ b \\
&= (-1 + q + 1 - q + q^2 - q^3 - q^2 + q^3)(a \circ b) \circ c \\
&\quad + (1 - q - 1 + q - q^2 + q^3 + q^2 - q^3)(a \circ c) \circ b \\
&\quad + (-q - q^2 + q + q^2)(b \circ c) \circ a + (-q^2 - q + q^2 + q)(c \circ b) \circ a \\
&= 0.
\end{aligned}$$

Lemma 4.2. *If $q \neq -2$, then*

$$\text{Alt}(\text{lei}^{(q)}) = -(q+2)(q-1) \text{alia}^{\left(\frac{-(2q+1)}{q+2}\right)}.$$

If $q = -2$, then

$$\text{Alt}(\text{lei}^{(-2)}) = 9 \text{ralia}.$$

Proof. Consider the case $q \neq -2$. We have

$$\begin{aligned}
&\text{lei}^{(q)}(t_1, t_2, t_3) + \text{lei}^{(q)}(t_2, t_3, t_1) + \text{lei}^{(q)}(t_3, t_1, t_2) - \text{lei}^{(q)}(t_2, t_1, t_3) - \text{lei}^{(q)}(t_3, t_2, t_1) \\
&\quad - \text{lei}^{(q)}(t_1, t_3, t_2) = (q-1)\{(2q+1)(t_1[t_2, t_3] + t_2[t_3, t_1]] + t_3[t_1, t_2]\} \\
&\quad - (q+2)([t_1, t_2]t_3 + [t_3, t_1]t_2 + [t_2, t_3]t_1) = (2-q-q^2) \text{ralia}^{\left(\frac{-(2q+1)}{q+2}\right)}.
\end{aligned}$$

The case $q = -2$ is considered in a similar manner. \square

Lemma 4.3. *Let L be a free Leibniz algebra with 3 generators, $q \in K, q \neq 0, \pm 1$. Then any multilinear identity of $L^{(q)}$ of degree 3 follows from the identities $\text{lei}^{(q)} = 0$ and $\text{lei}_1^{(q)} = 0$. If $q \neq -2$ then $\text{lei}_1^{(q)} = 0$ is a consequence of the identity $\text{lei}^{(q)} = 0$. If $q = -2$, then $\text{lei}^{(q)} = 0$ and $\text{lei}_1^{(q)} = 0$ are independent identities.*

Proof. Let $L = (L, \circ)$ be a free Leibniz algebra generated by three elements a, b, c . Write the q -commutator in $L^{(q)}$ by $uv = u \circ v + qv \circ u$.

The multilinear part of the free magma algebra (the algebra of non-commutative non-associative polynomials) in degree 3 has dimension 12. It is generated by the following 12 monomials:

$$\begin{aligned} e_1 &= e_1(t_1, t_2, t_3) = t_1(t_2t_3), & e_2 &= e_2(t_1, t_2, t_3) = t_2(t_3t_1), \\ e_3 &= e_3(t_1, t_2, t_3) = t_3(t_1t_2), & e_4 &= e_4(t_1, t_2, t_3) = t_2(t_1t_3), \\ e_5 &= e_5(t_1, t_2, t_3) = t_3(t_2t_1), & e_6 &= e_6(t_1, t_2, t_3) = t_1(t_3t_2), \\ e_7 &= e_7(t_1, t_2, t_3) = (t_1t_2)t_3, & e_8 &= e_8(t_1, t_2, t_3) = (t_2t_3)t_1, \\ e_9 &= e_9(t_1, t_2, t_3) = (t_3t_1)t_2, & e_{10} &= e_{10}(t_1, t_2, t_3) = (t_2t_1)t_3, \\ e_{11} &= e_{11}(t_1, t_2, t_3) = (t_3t_2)t_1, & e_{12} &= e_{12}(t_1, t_2, t_3) = (t_1t_3)t_2. \end{aligned}$$

Let $X = X(t_1, t_2, t_3) = \sum_{i=1}^{12} \lambda_i e_i(t_1, t_2, t_3)$ be a polynomial such that $X(a, b, c) = 0$ is an identity on $L^{(q)}$.

Substitute the generator elements $a, b, c \in L$ for the parameters t_1, t_2, t_3 . Write e_i instead of $e_i(a, b, c)$. We have

$$\begin{aligned} e_1 &= a \circ (b \circ c) + qa \circ (c \circ b) + q(b \circ c) \circ a + q^2(c \circ b) \circ a \\ &= (a \circ b) \circ c - (a \circ c) \circ b + q(a \circ c) \circ b - q(a \circ b) \circ c \\ &\quad + q(b \circ c) \circ a + q^2(c \circ b) \circ a. \end{aligned}$$

Similar calculations show that

$$\begin{aligned} e_2 &= (b \circ c) \circ a - (b \circ a) \circ c + q(b \circ a) \circ c - q(b \circ c) \circ a \\ &\quad + q(c \circ a) \circ b + q^2(a \circ c) \circ b, \end{aligned}$$

$$\begin{aligned} e_3 &= (c \circ a) \circ b - (c \circ b) \circ a + q(c \circ b) \circ a - q(c \circ a) \circ b \\ &\quad + q(a \circ b) \circ c + q^2(b \circ a) \circ c, \end{aligned}$$

$$\begin{aligned} e_4 &= (b \circ a) \circ c - (b \circ c) \circ a + q(b \circ c) \circ a - q(b \circ a) \circ c \\ &\quad + q(a \circ c) \circ b + q^2(c \circ a) \circ b, \end{aligned}$$

$$e_5 = (c \circ b) \circ a - (c \circ a) \circ b + q(c \circ a) \circ b - q(c \circ b) \circ a \\ + q(b \circ a) \circ c + q^2(a \circ b) \circ c,$$

$$e_6 = (a \circ c) \circ b - (a \circ b) \circ c + q(a \circ b) \circ c - q(a \circ c) \circ b \\ + q(c \circ b) \circ a + q^2(b \circ c) \circ a,$$

$$e_7 = (a \circ b) \circ c + q(b \circ a) \circ c + q(c \circ a) \circ b - q(c \circ b) \circ a \\ + q^2(c \circ b) \circ a - q^2(c \circ a) \circ b,$$

$$e_8 = (b \circ c) \circ a + q(c \circ b) \circ a + q(a \circ b) \circ c - q(a \circ c) \circ b \\ + q^2(a \circ c) \circ b - q^2(a \circ b) \circ c,$$

$$e_9 = (c \circ a) \circ b + q(a \circ c) \circ b + q(b \circ c) \circ a - q(b \circ a) \circ c \\ + q^2(b \circ a) \circ c - q^2(b \circ c) \circ a,$$

$$e_{10} = (b \circ a) \circ c + q(a \circ b) \circ c + q(c \circ b) \circ a - q(c \circ a) \circ b \\ + q^2(c \circ a) \circ b - q^2(c \circ b) \circ a,$$

$$e_{11} = (c \circ b) \circ a + q(b \circ c) \circ a + q(a \circ c) \circ b - q(a \circ b) \circ c \\ + q^2(a \circ b) \circ c - q^2(a \circ c) \circ b,$$

$$e_{12} = (a \circ c) \circ b + q(c \circ a) \circ b + q(b \circ a) \circ c - q(b \circ c) \circ a \\ + q^2(b \circ c) \circ a - q^2(b \circ a) \circ c.$$

So,

$$X =$$

$$(\lambda_1 - q\lambda_1 + q\lambda_3 + q^2 \lambda_5 - \lambda_6 + q\lambda_6 + \lambda_7 + q\lambda_8 - q^2 \lambda_8 + q\lambda_{10} - q\lambda_{11} + q^2 \lambda_{11})(a \circ b) \circ c \\ + (-\lambda_1 + q\lambda_1 + q^2 \lambda_2 + q\lambda_4 + \lambda_6 - q\lambda_6 - q\lambda_8 + q^2 \lambda_8 + q\lambda_9 + q\lambda_{11} - q^2 \lambda_{11} + \lambda_{12})(a \circ c) \circ b \\ + (-\lambda_2 + q\lambda_2 + q^2 \lambda_3 + \lambda_4 - q\lambda_4 + q\lambda_5 + q\lambda_7 - q\lambda_9 + q^2 \lambda_9 + \lambda_{10} + q\lambda_{12} - q^2 \lambda_{12})(b \circ a) \circ c \\ + (q\lambda_1 + \lambda_2 - q\lambda_2 - \lambda_4 + q\lambda_4 + q^2 \lambda_6 + \lambda_8 + q\lambda_9 - q^2 \lambda_9 + q\lambda_{11} - q\lambda_{12} + q^2 \lambda_{12})(b \circ c) \circ a \\ + (q\lambda_2 + \lambda_3 - q\lambda_3 + q^2 \lambda_4 - \lambda_5 + q\lambda_5 + q\lambda_7 - q^2 \lambda_7 + \lambda_9 - q\lambda_{10} + q^2 \lambda_{10} + q\lambda_{12})(c \circ a) \circ b \\ + (q^2 \lambda_1 - \lambda_3 + q\lambda_3 + \lambda_5 - q\lambda_5 + q\lambda_6 - q\lambda_7 + q^2 \lambda_7 + q\lambda_8 + q\lambda_{10} - q^2 \lambda_{10} + \lambda_{11})(c \circ b) \circ a.$$

Thus we obtain the following system of equations

$$\begin{aligned}
 (1 - q)\lambda_1 + q\lambda_3 + q^2 \lambda_5 + (q - 1)\lambda_6 + \lambda_7 + (q - q^2)\lambda_8 + q\lambda_{10} + (q^2 - q)\lambda_{11} &= 0, \\
 (q - 1)\lambda_1 + q^2\lambda_2 + q\lambda_4 + (1 - q)\lambda_6 + (q^2 - q)\lambda_8 + q\lambda_9 + (q - q^2)\lambda_{11} + \lambda_{12} &= 0, \\
 (q - 1)\lambda_2 + q^2\lambda_3 + (1 - q)\lambda_4 + q\lambda_5 + q\lambda_7 + (q^2 - q)\lambda_9 + \lambda_{10} + (q - q^2)\lambda_{12} &= 0, \\
 q\lambda_1 + (1 - q)\lambda_2 + (q - 1)\lambda_4 + q^2\lambda_6 + \lambda_8 + (q - q^2)\lambda_9 + q\lambda_{11} + (q^2 - q)\lambda_{12} &= 0, \\
 q\lambda_2 + (1 - q)\lambda_3 + q^2\lambda_4 + (q - 1)\lambda_5 + (q - q^2)\lambda_7 + \lambda_9 + (q^2 - q)\lambda_{10} + q\lambda_{12} &= 0, \\
 q^2\lambda_1 + (q - 1)\lambda_3 + (1 - q)\lambda_5 + q\lambda_6 + (q^2 - q)\lambda_7 + q\lambda_8 + (q - q^2)\lambda_{10} + \lambda_{11} &= 0.
 \end{aligned}$$

The 6×6 -determinant composed of the first 6 rows is $(1 - q)^5 q^3 (1 + q)^3 (q + 2)$. So, this system has rank 6 if $q^2 \neq 1, q \neq 0, -2$. One can choose $\lambda_i, 7 \leq i \leq 12$, as free parameters. Now, we consider two cases.

Suppose that $q \neq -2$. In this case the system has the following solution

$$\begin{aligned}
 \lambda_1 &= -\frac{-1 + q + q^2}{(q + 2)q}(\lambda_7 + \lambda_8 + \lambda_9 + (1 - q - q^2)\lambda_{10} + (1 + q)\lambda_{11} - \lambda_{12}), \\
 \lambda_2 &= -\frac{1}{(q + 2)q}(\lambda_7 + (q^2 + q - 1)\lambda_8 + \lambda_9 - \lambda_{10} + (1 - q - q^2)\lambda_{11} + (q + 1)\lambda_{12}), \\
 \lambda_3 &= -\frac{1}{(q + 2)q}(\lambda_7 + \lambda_8 + (q^2 + q - 1)\lambda_9 + (q + 1)\lambda_{10} - \lambda_{11} - (q^2 + q - 1)\lambda_{12}), \\
 \lambda_4 &= -\frac{1}{(q + 2)q}((1 - q - q^2)\lambda_7 - \lambda_8 + (q + 1)\lambda_9 + (q^2 + q - 1)\lambda_{10} + \lambda_{11} + \lambda_{12}), \\
 \lambda_5 &= -\frac{1}{(q + 2)q}((1 + q)\lambda_7 - (q^2 + q - 1)\lambda_8 - \lambda_9 + \lambda_{10} + (q^2 + q - 1)\lambda_{11} + \lambda_{12}), \\
 \lambda_6 &= -\frac{1}{(q + 2)q}(-\lambda_7 + (q + 1)\lambda_8 + (1 - q - q^2)\lambda_9 + \lambda_{10} + \lambda_{11} + (q^2 + q - 1)\lambda_{12}).
 \end{aligned}$$

Substitute these expressions for $\lambda_i, 1 \leq i \leq 6$, in $X(t_1, t_2, t_3)$ and collect the coefficients of $\lambda_j, 7 \leq j \leq 12$. We obtain a presentation of the polynomial $X(t_1, t_2, t_3)$ as a linear combination of the following 6 polynomials

$$\begin{aligned}
 f_1 &= (q - 1)t_1(t_2t_3) - (q^3 - q + 1)t_1(t_3t_2) - (q - 1)t_2(t_1t_3) \\
 &\quad - (q^2 + q - 1)t_2(t_3t_1) + (q^3 - q)t_3(t_1t_2) + (q^3 + q^2 - q)(t_1t_3)t_2 + q(t_2t_3)t_1,
 \end{aligned}$$

$$f_2 = (-1 + q^2)t_1(t_2t_3) - t_1(t_3t_2) - (q^2 - 1)t_2(t_1t_3) - qt_3(t_1t_2) \\ + (q^2 + q - 1)(t_2t_1)t_3 + (t_2t_3)t_1,$$

$$f_3 = (-q^3 + q - 1)t_1(t_2t_3) - t_1(t_3t_2) + (q^3 - q + 1)t_2(t_1t_3) \\ - (q^2 + q - 1)t_2(t_3t_1 - qt_3(t_1t_2)) + (q^3 + q^2 - q)(t_1t_2)t_3 + (q^2 + q)(t_2t_3)t_1,$$

$$f_4 = -t_1(t_2t_3) - (1 + q)t_1(t_3t_2) + t_2(t_1t_3) - (q^2 + q - 1)t_2(t_3t_1) \\ - t_3(t_1t_2) + (q^2 + q - 1)t_3(t_2t_1) + (q^2 + 2q)(t_2t_3)t_1,$$

$$f_5 = (1 - q)t_1(t_2t_3) + (q^3 - q + 1)t_1(t_3t_2) - q^2t_2(t_1t_3) - (q^3 - q)t_3(t_1t_2) \\ - q(t_2t_3)t_1 + (q^3 + q^2 - q)(t_3t_1)t_2,$$

$$f_6 = -t_1(t_2t_3) - t_1(t_3t_2) + q(t_2t_3)t_1 + q((t_3t_2)t_1).$$

We see that if $q^2 \neq 1, q \neq -2$, then

$$f_1 = \frac{1}{(q-1)(q+1)(q+2)}(-\text{lei}^{(q)}(t_1, t_2, t_3) - (-1 + q + q^2)\text{lei}^{(q)}(t_2, t_1, t_3) \\ + (-1 + q + q^2)^2\text{lei}^{(q)}(t_3, t_1, t_2) + (-1 + q + q^2)\text{lei}^{(q)}(t_3, t_2, t_1)),$$

$$f_2 = \text{lei}^{(q)},$$

$$f_3 = \frac{1}{(q-1)(q+1)(q+2)}(-(1 + q)\text{lei}^{(q)}(t_1, t_2, t_3) \\ + (-1 + q + q^2)^2\text{lei}^{(q)}(t_2, t_1, t_3) - (-1 + q + q^2)\text{lei}^{(q)}(t_3, t_1, t_2) \\ + (1 + q)(-1 + q + q^2)\text{lei}^{(q)}(t_3, t_2, t_1)),$$

$$f_4 = \frac{1}{(q+1)(q-1)}(-\text{lei}^{(q)}(t_1, t_2, t_3) + (-1 + q + q^2)\text{lei}^{(q)}(t_3, t_2, t_1)),$$

$$f_5 = \frac{1}{(q-1)(q+1)(q+2)}(\text{lei}^{(q)}(t_1, t_2, t_3) + (-1 + q + q^2)^2\text{lei}^{(q)}(t_1, t_3, t_2) \\ - (-1 + q + q^2)\text{lei}^{(q)}(t_2, t_3, t_1) - (-1 + q + q^2)\text{lei}^{(q)}(t_3, t_2, t_1)),$$

$$f_6 = \frac{1}{(q-1)(q+1)(q+2)}(-\text{lei}^{(q)}(t_1, t_2, t_3) - \text{lei}^{(q)}(t_1, t_3, t_2) \\ + (-1 + q + q^2)\text{lei}^{(q)}(t_2, t_3, t_1) + (-1 + q + q^2)\text{lei}^{(q)}(t_3, t_2, t_1)).$$

Now, we consider the case $q = -2$. In this case, similar arguments show

that X is a linear combination of the following polynomials

$$g_1 = t_3(t_2t_1) + 2/3(t_1t_2)t_3 + 4/3(t_1t_3)t_2 + 4/3(t_2t_1)t_3 - 4/3(t_2t_3)t_1 + 5/3(t_3t_1)t_2 - 5/3(t_3t_2)t_1,$$

$$g_2 = t_2(t_3t_1) + 4/3(t_1t_2)t_3 + 2/3(t_1t_3)t_2 + 5/3(t_2t_1)t_3 - 5/3(t_2t_3)t_1 + 4/3(t_3t_1)t_2 - 4/3(t_3t_2)t_1,$$

$$g_3 = t_1(t_2t_3) - 5/3(t_1t_2)t_3 + 5/3(t_1t_3)t_2 - 4/3(t_2t_1)t_3 + 4/3(t_2t_3)t_1 + 4/3(t_3t_1)t_2 + 2/3(t_3t_2)t_1,$$

$$g_4 = t_1(t_3t_2) + 5/3(t_1t_2)t_3 - 5/3(t_1t_3)t_2 + 4/3(t_2t_1)t_3 + 2/3(t_2t_3)t_1 - 4/3(t_3t_1)t_2 + 4/3(t_3t_2)t_1,$$

$$g_5 = t_2(t_1t_3) - 4/3(t_1t_2)t_3 + 4/3(t_1t_3)t_2 - 5/3(t_2t_1)t_3 + 5/3(t_2t_3)t_1 + 2/3(t_3t_1)t_2 + 4/3(t_3t_2)t_1,$$

$$g_6 = t_3(t_1t_2 + 4/3(t_1t_2)t_3 - 4/3(t_1t_3)t_2 + 2/3(t_2t_1)t_3 + 4/3(t_2t_3)t_1 - 5/3(t_3t_1)t_2 + 5/3(t_3t_2)t_1).$$

We have

$$g_1 - 1/3(4 \text{lei}^{(-2)}(t_1, t_2, t_3) + 3 \text{lei}^{(-2)}(t_1, t_3, t_2) + 2 \text{lei}^{(-2)}(t_2, t_1, t_3) + 4 \text{lei}_1^{(-2)}(t_1, t_2, t_3) - \text{lei}_1^{(-2)}(t_2, t_1, t_3)) = 1/3 \text{ralia}(t_1, t_2, t_3),$$

$$g_2 - 1/3(5 \text{lei}^{(-2)}(t_1, t_2, t_3) + 6 \text{lei}^{(-2)}(t_1, t_3, t_2) + 4 \text{lei}^{(-2)}(t_2, t_1, t_3) + 5 \text{lei}_1^{(-2)}(t_1, t_2, t_3) + \text{lei}_1^{(-2)}(t_2, t_1, t_3)) = 8/3 \text{ralia}(t_1, t_2, t_3),$$

$$g_3 - 1/3(-4 \text{lei}^{(-2)}(t_1, t_2, t_3) - 6 \text{lei}^{(-2)}(t_1, t_3, t_2) - 5 \text{lei}^{(-2)}(t_2, t_1, t_3) - 4 \text{lei}_1^{(-2)}(t_1, t_2, t_3) - 5 \text{lei}_1^{(-2)}(t_2, t_1, t_3)) = -10/3 \text{ralia}(t_1, t_2, t_3),$$

$$g_4 - 1/3(4 \text{lei}^{(-2)}(t_1, t_2, t_3) + 6 \text{lei}^{(-2)}(t_1, t_3, t_2) + 5 \text{lei}^{(-2)}(t_2, t_1, t_3) + \text{lei}_1^{(-2)}(t_1, t_2, t_3) + 5 \text{lei}_1^{(-2)}(t_2, t_1, t_3)) = 10/3 \text{ralia}(t_1, t_2, t_3),$$

$$g_5 - 1/3(-5 \text{lei}^{(-2)}(t_1, t_2, t_3) - 6 \text{lei}^{(-2)}(t_1, t_3, t_2) - 4 \text{lei}^{(-2)}(t_2, t_1, t_3) - 5 \text{lei}_1^{(-2)}(t_1, t_2, t_3) - 4 \text{lei}_1^{(-2)}(t_2, t_1, t_3)) = -8/3 \text{ralia}(t_1, t_2, t_3),$$

$$g_6 - 1/3(2\text{lei}^{(-2)}(t_1, t_2, t_3) + 3\text{lei}^{(-2)}(t_1, t_3, t_2) + 4\text{lei}^{(-2)}(t_2, t_1, t_3) - \text{lei}_1^{(-2)}(t_1, t_2, t_3) + 4\text{lei}_1^{(-2)}(t_2, t_1, t_3)) = 8/3 \text{ralia}(t_1, t_2, t_3).$$

By Lemma 4.2 $\text{ralia} = 0$ is a consequence of the identity $\text{lei}^{(q)} = 0$. Therefore, all the identities $g_i = 0$ are consequences of the identities $\text{lei}^{(q)} = 0$ and $\text{lei}_1^{(q)} = 0$.

We have proved that any identity of degree 3 of $L^{(q)}$ for $q = -2$ follows from the identities $\text{lei}^{(q)} = 0$ and $\text{lei}_1^{(q)} = 0$. Notice that the equation

$$\begin{aligned} \text{lei}_1^{(q)}(a, b, c) &= \mu_1 \text{lei}^{(q)}(a, b, c) + \mu_2 \text{lei}^{(q)}(b, c, a) + \mu_3 \text{lei}^{(q)}(c, a, b) \\ &\quad + \mu_4 \text{lei}^{(q)}(b, a, c) + \mu_5 \text{lei}^{(q)}(c, b, a) + \mu_6 \text{lei}^{(q)}(a, c, b) \end{aligned}$$

in $L^{(q)}$ with unknowns $\mu_1, \mu_2, \mu_3, \mu_4, \mu_5, \mu_6$ is not solvable. Therefore, this system of identities $\text{lei}^{(q)} = 0, \text{lei}_1^{(q)} = 0$ is $\mathfrak{L}\text{ei}^{(q)}$ -minimal if $q = -2$.

Lemma 4.4. *Suppose that $q \neq 0, \pm 1$ and an algebra (A, \star) satisfies the identities $\text{lei}^{(q)} = 0$ and $\text{lei}_1^{(q)} = 0$. Then the algebra (A, \circ) , where $a \circ b = (1 - q^2)^{-1}(a \star b - q b \star a)$, is a (right)-Leibniz algebra, and the algebras (A, \star) and (A, \circ_q) are isomorphic.*

Proof. One checks that

$$\begin{aligned} \text{lei}(t_1, t_2, t_3) &= -2 \text{lei}^{(q)}(t_1, t_2, t_3) - 2/3 \text{lei}^{(q)}(t_1, t_3, t_2) - \text{lei}^{(q)}(t_2, t_1, t_3) \\ &\quad + 2/3 (\text{lei}^{(q)}(t_2, t_3, t_1) - 2\text{lei}_1^{(q)}(t_1, t_2, t_3) - \text{lei}_1^{(q)}(t_2, t_1, t_3)) \end{aligned}$$

for $q^2 \neq 1, q = -2$, and

$$\begin{aligned} \text{lei}(t_1, t_2, t_3) &= \frac{1}{(q^2 - 1)(q + 2)} (q(q + 1) \text{lei}^{(q)}(t_1, t_2, t_3) \\ &\quad - (-1 + 2q + q^2) \text{lei}^{(q)}(t_1, t_3, t_2) - (q + 1) \text{lei}^{(q)}(t_2, t_1, t_3) \\ &\quad + (1 - q + q^2 + q^3) \text{lei}^{(q)}(t_2, t_3, t_1) + (q + 1) \text{lei}^{(q)}(t_3, t_1, t_2) \\ &\quad - (q + q^2) \text{lei}^{(q)}(t_3, t_2, t_1)) \end{aligned}$$

for $q^2 \neq 1, q \neq -2$.

Therefore, for any algebra (A, \star) with identities $\text{lei}^{(q)} = 0$ and $\text{lei}_1^{(q)} = 0$ the algebra (A, \circ) , where $a \circ b = (1 - q^2)^{-1}(a \star b - q b \star a)$, satisfies the identity $\text{lei} = 0$. It is evident that

$$\begin{aligned} a \circ_q b &= (1 - q^2)^{-1}(a \circ b + q b \circ a) \\ &= (1 - q^2)^{-1}(a \star b - q b \star a + q b \star a - q^2 a \star b) \\ &= a \star b. \end{aligned}$$

Proof of Theorems 1.1 and 1.2. By Lemmas 4.1, 4.3, 4.4 our theorems are true. \square

5. Leibniz-Lie algebras. In this section we study identities for Leibniz-Lie algebras, i.e., algebras $(A, [,])$ under -1 -commutator for Leibniz algebras (A, \circ) .

Note that $lei_1(t_1, t_2, t_3, t_4, t_5)$ has type $(3, 2)$, i.e., it is skew-symmetric in t_1, t_2, t_3 and in t_4, t_5 , and $lei_2(t_1, t_2, t_3, t_4, t_5)$ has type $(1, 4)$, is skew-symmetric in t_2, t_3, t_4, t_5 .

Let

$$\begin{aligned}
 lei_3^{(-1)}(t_1, t_2, t_3, t_4, t_5) = & ((t_1t_2)t_3)(t_4t_5) + ((t_1t_2)t_5)(t_3t_4) + ((t_1t_2)(t_3t_4))t_5 - 2((t_1t_2)(t_3t_5))t_4 \\
 & + ((t_1t_2)(t_4t_5))t_3 - ((t_1t_3)t_5)(t_2t_4) + ((t_1t_3)(t_2t_4))t_5 - ((t_1t_4)t_3)(t_2t_5) \\
 & + ((t_1t_4)t_5)(t_2t_3) + ((t_1t_4)(t_2t_3))t_5 - ((t_1t_4)(t_2t_5))t_3 + 2((t_1t_4)(t_3t_5))t_2 \\
 & + ((t_1t_5)t_3)(t_2t_4) - ((t_1t_5)(t_2t_4))t_3 + ((t_2t_3)t_5)(t_1t_4) + ((t_2t_4)t_3)(t_1t_5) \\
 & - ((t_2t_4)t_5)(t_1t_3) - 2((t_2t_4)(t_3t_5))t_1 - ((t_2t_5)t_3)(t_1t_4) + ((t_3t_5)t_4)(t_1t_2) \\
 & + 2(((t_1t_2)t_3)t_5)t_4 - 6(((t_1t_2)t_4)t_3)t_5 + 6(((t_1t_2)t_4)t_5)t_3 - 2(((t_1t_2)t_5)t_3)t_4 \\
 & - 2(((t_1t_3)t_2)t_4)t_5 + 2(((t_1t_3)t_4)t_2)t_5 + 6(((t_1t_3)t_5)t_2)t_4 - 6(((t_1t_3)t_5)t_4)t_2 \\
 & + 6(((t_1t_4)t_2)t_3)t_5 - 6(((t_1t_4)t_2)t_5)t_3 - 2(((t_1t_4)t_3)t_5)t_2 + 2(((t_1t_4)t_5)t_3)t_2 \\
 & + 2(((t_1t_5)t_2)t_4)t_3 - 6(((t_1t_5)t_3)t_2)t_4 + 6(((t_1t_5)t_3)t_4)t_2 - 2(((t_1t_5)t_4)t_2)t_3 \\
 & + 2(((t_2t_3)t_1)t_4)t_5 - 2(((t_2t_3)t_4)t_1)t_5 - 6(((t_2t_3)t_5)t_1)t_4 + 6(((t_2t_3)t_5)t_4)t_1 \\
 & - 6(((t_2t_4)t_1)t_3)t_5 + 6(((t_2t_4)t_1)t_5)t_3 + 2(((t_2t_4)t_3)t_5)t_1 - 2(((t_2t_4)t_5)t_3)t_1 \\
 & - 2(((t_2t_5)t_1)t_4)t_3 + 6(((t_2t_5)t_3)t_1)t_4 - 6(((t_2t_5)t_3)t_4)t_1 + 2(((t_2t_5)t_4)t_1)t_3 \\
 & + 2(((t_3t_4)t_1)t_2)t_5 - 2(((t_3t_4)t_2)t_1)t_5 - 4(((t_3t_4)t_5)t_1)t_2 + 4(((t_3t_4)t_5)t_2)t_1 \\
 & + 4(((t_3t_5)t_1)t_2)t_4 - 4(((t_3t_5)t_1)t_4)t_2 - 4(((t_3t_5)t_2)t_1)t_4 + 4(((t_3t_5)t_2)t_4)t_1 \\
 & + 2(((t_3t_5)t_4)t_1)t_2 - 2(((t_3t_5)t_4)t_2)t_1 + 2(((t_4t_5)t_1)t_2)t_3 - 2(((t_4t_5)t_2)t_1)t_3 \\
 & - 4(((t_4t_5)t_3)t_1)t_2 + 4(((t_4t_5)t_3)t_2)t_1.
 \end{aligned}$$

Statements below need long calculations. We omit them. Details one can find in our preprint [5].

Lemma 5.1. *The identity $\text{lei}_3^{(-1)} = 0$ is a consequence of the identities $\text{leilie}_1 = 0$, $\text{leilie}_2 = 0$ and the anti-commutativity identity.*

Lemma 5.2. *Let (A, \circ) be a Leibniz algebra. Then the Leibniz-Lie algebra $(A, [,])$ satisfies the identities $\text{lei}_1^{(-1)} = 0$, $\text{leilie}_2 = 0$.*

Lemma 5.3. *Any identity of degree 4 for $\mathfrak{Lci}^{(-1)}$ follows from the identity $\text{acom} = 0$.*

Proof. Let, working modulo the identity of anti-commutativity

$$\begin{aligned} X_4(t_1, t_2, t_3, t_4) = & \lambda_1(t_1t_2)(t_3t_4) + \lambda_2(t_1t_3)(t_2t_4) + \lambda_3(t_2t_3)(t_1t_4) + \lambda_4((t_1t_2)t_3)t_4 \\ & + \lambda_{10}((t_1t_2)t_4)t_3 + \lambda_5((t_1t_3)t_2)t_4 + \lambda_{11}((t_1t_3)t_4)t_2 + \lambda_6((t_1t_4)t_2)t_3 \\ & + \lambda_{12}((t_1t_4)t_3)t_2 + \lambda_7((t_2t_3)t_1)t_4 + \lambda_{13}((t_2t_3)t_4)t_1 + \lambda_8((t_2t_4)t_1)t_3 \\ & + \lambda_{14}((t_2t_4)t_3)t_1 + \lambda_9((t_3t_4)t_1)t_2 + \lambda_{15}((t_3t_4)t_2)t_1 \end{aligned}$$

be a generic multilinear polynomial of degree 4. For t_1, t_2, t_3, t_4 , we substitute the elements a, b, c, d of the free Leibniz algebra, and calculate $X_4(a, b, c, d)$ under the commutator $[u, v] = u \circ v - v \circ u$. We obtain

$$\begin{aligned} X_4(a, b, c, d) = & (2\lambda_1 + \lambda_4 - 2\lambda_7 - 4\lambda_{13} + 4\lambda_{15})(((a \circ b) \circ c) \circ d) \\ & + (-2\lambda_1 - 2\lambda_8 + \lambda_{10} - 4\lambda_{14} - 4\lambda_{15})(((a \circ b) \circ d) \circ c) \\ & + (2\lambda_2 + \lambda_5 + 2\lambda_7 + 4\lambda_{13} + 4\lambda_{14})(((a \circ c) \circ b) \circ d) \\ & + (-2\lambda_2 - 2\lambda_9 + \lambda_{11} - 4\lambda_{14} - 4\lambda_{15})(((a \circ c) \circ d) \circ b) \\ & + (-2\lambda_3 + \lambda_6 + 2\lambda_8 + 4\lambda_{13} + 4\lambda_{14})(((a \circ d) \circ b) \circ c) \\ & + (2\lambda_3 + 2\lambda_9 + \lambda_{12} - 4\lambda_{13} + 4\lambda_{15})(((a \circ d) \circ c) \circ b) \\ & + (-2\lambda_1 - \lambda_4 - 2\lambda_5 + 4\lambda_9 - 4\lambda_{11})(((b \circ a) \circ c) \circ d) \\ & + (2\lambda_1 - 2\lambda_6 - 4\lambda_9 - \lambda_{10} - 4\lambda_{12})(((b \circ a) \circ d) \circ c) \\ & + (2\lambda_3 + 2\lambda_5 + \lambda_7 + 4\lambda_{11} + 4\lambda_{12})(((b \circ c) \circ a) \circ d) \\ & + (-2\lambda_3 - 4\lambda_9 - 4\lambda_{12} + \lambda_{13} - 2\lambda_{15})(((b \circ c) \circ d) \circ a) \\ & + (-2\lambda_2 + 2\lambda_6 + \lambda_8 + 4\lambda_{11} + 4\lambda_{12})(((b \circ d) \circ a) \circ c) \end{aligned}$$

$$\begin{aligned}
 &+ (2\lambda_2 + 4\lambda_9 - 4\lambda_{11} + \lambda_{14} + 2\lambda_{15})(((b \circ d) \circ c) \circ a) \\
 &+ (-2\lambda_2 - 2\lambda_4 - \lambda_5 + 4\lambda_8 - 4\lambda_{10})(((c \circ a) \circ b) \circ d) \\
 &+ (2\lambda_2 - 4\lambda_6 - 4\lambda_8 - \lambda_{11} - 2\lambda_{12})(((c \circ a) \circ d) \circ b) \\
 &+ (-2\lambda_3 + 2\lambda_4 + 4\lambda_6 - \lambda_7 + 4\lambda_{10})(((c \circ b) \circ a) \circ d) \\
 &+ (2\lambda_3 - 4\lambda_6 - 4\lambda_8 - \lambda_{13} - 2\lambda_{14})(((c \circ b) \circ d) \circ a) \\
 &+ (-2\lambda_1 + 4\lambda_6 + \lambda_9 + 4\lambda_{10} + 2\lambda_{12})(((c \circ d) \circ a) \circ b) \\
 &+ (2\lambda_1 + 4\lambda_8 - 4\lambda_{10} + 2\lambda_{14} + \lambda_{15})(((c \circ d) \circ b) \circ a) \\
 &+ (2\lambda_3 - 4\lambda_4 - \lambda_6 + 4\lambda_7 - 2\lambda_{10})(((d \circ a) \circ b) \circ c) \\
 &+ (-2\lambda_3 - 4\lambda_5 - 4\lambda_7 - 2\lambda_{11} - \lambda_{12})(((d \circ a) \circ c) \circ b) \\
 &+ (2\lambda_2 + 4\lambda_4 + 4\lambda_5 - \lambda_8 + 2\lambda_{10})(((d \circ b) \circ a) \circ c) \\
 &+ (-2\lambda_2 - 4\lambda_5 - 4\lambda_7 - 2\lambda_{13} - \lambda_{14})(((d \circ b) \circ c) \circ a) \\
 &+ (2\lambda_1 + 4\lambda_4 + 4\lambda_5 - \lambda_9 + 2\lambda_{11})(((d \circ c) \circ a) \circ b) \\
 &+ (-2\lambda_1 - 4\lambda_4 + 4\lambda_7 + 2\lambda_{13} - \lambda_{15})(((d \circ c) \circ b) \circ a).
 \end{aligned}$$

Since all 24 left-bracketed elements like $((a \circ b) \circ c) \circ d$ are linear independent elements, the condition $X_4(a, b, c, d) = 0$ gives us the system of 24 linear equations in 15 unknowns $\lambda_i, i = 1, \dots, 15$. We see that the rank of this system is 15 and our system has the trivial solution only: $\lambda_i = 0$ for all $i = 1, 2, \dots, 15$. In other words, any multilinear identity of degree 4 for $\mathfrak{L}\mathfrak{e}\mathfrak{i}^{(-1)}$ follows from the identity $\text{acom} = 0$.

Lemma 5.4. *Any identity of degree 5 for the free Leibniz algebra follows from the identities $\text{lelie}_1 = 0, \text{lelie}_2 = 0, \text{lei}_3^{(-1)} = 0$.*

Proof. Let $f = f(t_1, \dots, t_5)$ be a non-commutative non-associative polynomial such that $f = 0$ is an identity for any right-Leibniz algebra. Notice that there exist 105 anti-commutative non-associative polynomials. Present f as a linear combination of these 105 elements.

Insert in f the elements of the free Leibniz algebra generated by 5 elements u_1, u_2, u_3, u_4, u_5 and calculate the polynomial f under the commutator $[u, v] = u \circ v - v \circ u$, where $(u, v) \mapsto u \circ v$ is the multiplication in a free (right)-Leibniz algebra. Expand this expression in terms of the multiplication \circ using the Leibniz rule

$$u \circ (v \circ w) = (u \circ v) \circ w - (u \circ w) \circ v.$$

We obtain an element which is a linear combination of 120 elements of the form $((u_{\sigma(1)} \circ u_{\sigma(2)}) \circ u_{\sigma(3)}) \circ u_{\sigma(4)} \circ u_{\sigma(5)}$, where $\sigma \in \text{Sym}_5$. The identity condition $f = 0$ on $L^{(-1)}$ gives us 120 linear equations in 105 unknowns λ_i . Solve this system of equations. We do this using the computer system *Mathematica*. We find out that the system has 14 free parameters. It shows that f can be presented as a linear combination of the 14 polynomials given below

$$f_1 = \text{leilie}_1,$$

$$f_2 = \text{leilie}_1(t_1, t_2, t_4, t_3, t_5),$$

$$f_3 = \text{leilie}_1(t_1, t_2, t_5, t_3, t_4),$$

$$f_4 = \text{leilie}_1(t_1, t_3, t_4, t_2, t_5),$$

$$f_5 = \text{leilie}_1(t_2, t_3, t_4, t_1, t_5),$$

$$f_6 = \text{leilie}_1(t_1, t_3, t_5, t_2, t_4),$$

$$f_7 = \text{leilie}_1(t_2, t_3, t_5, t_1, t_4),$$

$$f_9 = \text{leilie}_1(t_1, t_4, t_5, t_2, t_3) + \text{leilie}_2(t_1, t_2, t_3, t_4, t_5),$$

$$f_{10} = (\text{leilie}_1(t_3, t_4, t_5, t_1, t_2) + 2 \text{leilie}_2(t_1, t_2, t_3, t_4, t_5) + \text{lei}_3^{(-1)}(t_1, t_2, t_3, t_4, t_5) - \text{lei}_3^{(-1)}(t_1, t_2, t_3, t_5, t_4) - \text{lei}_3^{(-1)}(t_1, t_2, t_4, t_3, t_5))/2,$$

$$f_{11} = (2 \text{leilie}_1(t_2, t_4, t_5, t_1, t_3) + \text{leilie}_1(t_3, t_4, t_5, t_1, t_2) + 2 \text{leilie}_2(t_1, t_2, t_3, t_4, t_5) + \text{lei}_3^{(-1)}(t_1, t_2, t_3, t_4, t_5) - \text{lei}_3^{(-1)}(t_1, t_2, t_3, t_5, t_4) - \text{lei}_3^{(-1)}(t_1, t_2, t_4, t_3, t_5))/2,$$

$$f_{12} = (\text{leilie}_1(t_3, t_4, t_5, t_1, t_2) + \text{lei}_3^{(-1)}(t_1, t_2, t_3, t_4, t_5) + \text{lei}_3^{(-1)}(t_1, t_2, t_3, t_5, t_4) - \text{lei}_3^{(-1)}(t_1, t_2, t_4, t_3, t_5))/2,$$

$$f_{14} = (\text{leilie}_1(t_3, t_4, t_5, t_1, t_2) + \text{lei}_3^{(-1)}(t_1, t_2, t_3, t_4, t_5) - \text{lei}_3^{(-1)}(t_1, t_2, t_3, t_5, t_4) + \text{lei}_3^{(-1)}(t_1, t_2, t_4, t_3, t_5))/2.$$

So, by Lemma 5.1 the 9-term polynomial leilie_1 and the 60-term polynomial leilie_2 form a base of multilinear identities of degree 5.

Proof of Theorem 1.3. Follows from Lemmas 5.1, 5.2, 5.3 and 5.4. \square

6. Leibniz-Jordan algebras.

Proof of Theorem 1.6. It is easy to check that $\text{leijor} = 0$ is an identity for any algebra of the form $A^{(1)}$, where A is a Leibniz algebra.

Let A be an associative algebra and let M be a right module over A . Then $A^{(-1)}$ is a Lie algebra and M can be made into an antisymmetric $A^{(-1)}$ -module. Let $L = A + M$ be the standard Leibniz algebra corresponding to these Lie and antisymmetric module structures. If we denote by \star the multiplication in the Leibniz algebra L , then

$$(a + m) \star (b + n) = [a, b] + mb,$$

and

$$\{a + m, b + n\} = [a, b] + mb + [b, a] + na = na + mb.$$

In particular,

$$(3) \quad \{a, m\} = ma, \quad \{a, b\} = 0, \quad \{m, n\} = 0$$

for all $a, b \in A, m, n \in M$. Recall that

$$\{t_1, t_2\} = t_1t_2 + t_2t_1$$

is the Jordan commutator.

Suppose that $f = 0$ is a minimal identity for the Leibniz-Jordan algebra $(L, \{, \})$ which does not follow from the identity $\text{leijor} = 0$. We can assume that f is multilinear and $f = f(t_1, \dots, t_k)$ is a linear combination of left-bracketed monomials of the form $((t_{i_1}t_{i_2}) \cdots)t_{i_k}$. So,

$$f(t_1, \dots, t_k) = \sum_{\sigma \in \text{Sym}_k} \lambda_\sigma ((t_{\sigma(1)}t_{\sigma(2)}) \cdots)t_{\sigma(k)}$$

for some $\lambda_\sigma \in K$. Write the condition $f(a_1, \dots, a_{k-1}, m) = 0$ by using the multiplication rules (3) for Leibniz-Jordan algebras. We have

$$(4) \quad f(a_1, \dots, a_{k-1}, m) = \sum_{\sigma \in \text{Sym}_{k-1}} \lambda_\sigma ((ma_{\sigma(1)}) \cdots)a_{\sigma(k-1)}$$

for any $a_1, \dots, a_{k-1} \in A, m \in M$.

Take $A = \text{Mat}_n$ to be the matrix algebra and $M = K^n$ the n -dimensional natural module. Then conditions (4) imply that

$$\sum_{\sigma \in \text{Sym}_{k-1}} \lambda_\sigma ((a_{\sigma(1)}a_{\sigma(2)}) \cdots)a_{\sigma(k-1)} = 0$$

is an identity for Mat_n . By the Amitsur-Levitsky Theorem [1], matrix algebras have no identity of degree $k - 1$ if $k < 2n + 1$. So, $f = 0$ is not an identity for Leibniz-Jordan algebras of the form $\text{Mat}_n + K^n$ if $n > (k - 1)/2$. In other words, any s -identity for Leibniz-Jordan algebras follows from the identities $\text{leijor} = 0$, $\text{com} = 0$.

REFERENCES

- [1] S. A. AMITSUR, J. LEVITZKI. Minimal identities for algebras. *Proc. Amer. Math. Soc.* **1** (1950), 449–463.
- [2] A. BLOCH. On generalized notion of Lie algebras. *Dokl. Akad. Nauk SSSR* **165**, 3 (1965), 471–473.
- [3] A. S. DZHUMADIL'DAEV. q -chronological algebras. *Dokl. Akad. Nauk* **401**, 6 (2005), 731–732 (in Russian).
- [4] A. S. DZHUMADIL'DAEV. Zinbiel algebras under q -commutator. *Fund. Appl. Math.* **11**, 3 (2005), 57–78.
- [5] A. S. DZHUMADIL'DAEV. q -Leibniz algebras. Max-Planck Institute for Mathematics, Bonn, Preprint MPIM 2006-27.
- [6] M. K. KINYON, A. WEINSTEIN. Leibniz algebras, Courant algebroids, and multiplications on reductive homogeneous spaces. *Amer. J. Math.* **123**, 3 (2001), 525–550.
- [7] J.-L. LODAY, T. PIRASHVILI. Universal enveloping algebras of Leibniz algebras and (co)homology. *Math. Ann.* **296** (1993), 139–158.
- [8] J.-L. LODAY. Cup-product for Leibniz cohomology and dual Leibniz algebras. *Math. Scand.* **77**, 2 (1995), 189–196.
- [9] A. WEINSTEIN. Omni-Lie algebras. *RIMS Kokyuroku* **1176** (2000), 95–102; <http://xxx.lanl.gov/abs/math/9912190v2>.

Kazakh-British University

Tole bi 59, Almaty, 050000, Kazakhstan

e-mail: askar@math.kz

askar56@hotmail.com

Received September 22, 2006

Revised July 10, 2007