

Provided for non-commercial research and educational use.
Not for reproduction, distribution or commercial use.

Serdica

Mathematical Journal

Сердика

Математическо списание

The attached copy is furnished for non-commercial research and education use only.
Authors are permitted to post this version of the article to their personal websites or institutional repositories and to share with other researchers in the form of electronic reprints.
Other uses, including reproduction and distribution, or selling or licensing copies, or posting to third party websites are prohibited.

For further information on
Serdica Mathematical Journal
which is the new series of
Serdica Bulgaricae Mathematicae Publicationes
visit the website of the journal <http://www.math.bas.bg/~serdica>
or contact: Editorial Office
Serdica Mathematical Journal
Institute of Mathematics and Informatics
Bulgarian Academy of Sciences
Telephone: (+359-2)9792818, FAX:(+359-2)971-36-49
e-mail: serdica@math.bas.bg

REMARKS ON BLOW UP TIME FOR SOLUTIONS OF A NONLINEAR DIFFUSION SYSTEM WITH TIME DEPENDENT COEFFICIENTS

M. Marras

Communicated by T. Gramchev

ABSTRACT. We investigate the blow-up of the solutions to a nonlinear parabolic system with Robin boundary conditions and time dependent coefficients. We derive sufficient conditions on the nonlinearities and the initial data in order to obtain explicit lower and upper bounds for the blow up time t^* .

1. Introduction. The blow-up phenomena for solutions to linear and nonlinear parabolic equations and systems have been widely studied and different methods have been introduced in order to find bounds for blow-up time t^* . We refer to the books of Straughan [11] and Quittner-Souplet [10], and the papers of Vazquez [12] and Weissler [13], [14]. Recently Payne, Philippin, Schaefer in [5]–[6] and Payne, Philippin and Vernier-Piro in [7]–[8] (see also references therein) have considered nonlinear boundary value problems under different boundary

2010 *Mathematics Subject Classification*: 35K55, 35K60.

Key words: Parabolic problems, blow-up.

conditions, deriving lower and upper bounds for the blow-up time t^* as well as sufficient conditions for global existence of the solutions.

In this paper we investigate the blow-up phenomena of the classical solution $(u(x, t), v(x, t))$ of the following Robin system

$$(1.1) \quad \left\{ \begin{array}{l} \Delta u + K_1(t)f_1(v) = u_t \quad \text{in } \Omega \times (0, t^*), \\ \Delta v + K_2(t)f_2(u) = v_t \quad \text{in } \Omega \times (0, t^*), \\ \frac{\partial u}{\partial \nu}(x, t) = -\alpha u, \quad \text{on } \partial\Omega \times (0, t^*), \\ \frac{\partial v}{\partial \nu}(x, t) = -\beta v, \quad \text{on } \partial\Omega \times (0, t^*), \\ u(x, 0) = u_0(x) \geq 0, \quad \text{on } \Omega, \\ v(x, 0) = v_0(x) \geq 0, \quad \text{on } \Omega, \end{array} \right.$$

where Ω is a bounded domain in \mathbb{R}^N , $N \geq 2$ with $\partial\Omega$ sufficiently smooth, the coefficients $K_1(t)$, $K_2(t)$ and f_1 , f_2 positive functions and $\frac{\partial u}{\partial \nu}$, $\frac{\partial v}{\partial \nu}$ are the outward normal derivative of the vector-valued solution (u, v) on the boundary $\partial\Omega$, α and β are two positive constants satisfying

$$(1.2) \quad 0 \leq \alpha \leq \beta.$$

The initial data $u_0(x), v_0(x)$ are nonnegative smooth functions, such that $\frac{\partial u_0}{\partial \nu} = -\alpha u_0$, $\frac{\partial v_0}{\partial \nu} = -\beta v_0$, $x \in \partial\Omega$, while t^* stands for the blow-up time if blow up occurs, otherwise $t^* = \infty$. It is well known that the solution can fail to exist only by blowing up at finite time ([1], [2]) and the geometry of the domain Ω , the nonlinearities, the boundary data and the initial conditions greatly affect the evolution in time of the solution.

We note that u and v are non negative in x and $t \in (0, t^*)$ by the parabolic maximum principle applied to the the system (1.1).

The main aim of this paper is to derive explicit upper and lower bounds for the blow-up time for the solution of (1.1). We remark that if K_1, K_2 are constant, bounds for the blow-up time are derived in [3], where on $\partial\Omega \times (0, t^*)$ $\frac{\partial u}{\partial \nu} = g_1(u)$, $\frac{\partial v}{\partial \nu} = g_2(v)$, g_1, g_2 positive functions.

For systems with Dirichlet and Neumann boundary conditions we refer to [3] and [9].

We consider another class of parabolic systems with Robin type boundary conditions in [4], where sufficient conditions are introduced in order to obtain

global existence of the solution as well as upper and lower bounds for the blow-up, if blow-up occurs.

The paper is organized as follows. In Section 2 we derive a lower bound for the blow up time t^* under suitable conditions on data and for convex domain in \mathbb{R}^3 . In Section 3 we remove the restriction on the domain and under alternative conditions on data an upper bound is obtained.

Throughout the paper the notations $u_i = \frac{\partial u}{\partial x_i}$, $i = 1, \dots, N$ and $(\cdot)'$ for the derivative of the coefficients will be used and the summation convention over repeated indexes will be assumed.

2. Lower bound. Under suitable conditions on non linearities and Ω we get a differential inequality and we derive a lower bound for t^* .

To this end we will use a Sobolev-type inequality introduced in [3] which plays an important role in deriving the lower bound. However, it must be noted that the inequality holds only if we consider our domain $\Omega \subset \mathbb{R}^3$.

Let us suppose $K_1 \leq K_2$ and we define the auxiliary function

$$(2.1) \quad \Phi(t) = K_2^2(t) \int_{\Omega} (u^2 + v^2)^{2p} dx, \quad p > 1$$

We say that (u, v) blows up in Φ -measure when

$$\lim_{t \rightarrow t^*} \Phi(t) = \infty.$$

We now prove the following result

Theorem 2.1. *Let (u, v) be the solution of (1.1), in a convex domain $\Omega \subset \mathbb{R}^3$, blowing up at time t^* in Φ -measure. Assume that there exists a positive constant C_0 such that*

$$(2.2) \quad u f_1(v) + v f_2(u) \leq C_0 (u^2 + v^2)^{p+1}.$$

Moreover assume that $K_2(t)$ satisfies

$$(2.3) \quad \frac{K_2'(t)}{K_2(t)} \leq \delta, \quad \delta > 0.$$

Then the solution blows up in the Φ -measure and

$$(2.4) \quad t^* \geq T := \int_{\Phi(0)}^{\infty} \frac{d\eta}{\phi(\eta)},$$

where $\phi(\Phi) = \sum_{i=1}^3 (\xi_i \Phi^{h_i})$ with some if not all h_i greater than one.

Proof. We compute

$$\begin{aligned} \Phi'(t) &= 2 \left(\frac{K'_2}{K_2} \right) K_2^2 \int_{\Omega} (u^2 + v^2)^{2p} + 4pK_2^2 \int_{\Omega} (u^2 + v^2)^{2p-1} (u\Delta u + v\Delta v) dx \\ &\quad + 4pK_2^2 \int_{\Omega} (u^2 + v^2)^{2p-1} [K_1 u f_1(v) + K_2 v f_2(u)] dx. \end{aligned}$$

Now by (2.2), (2.3) and the assumption $K_1 \leq K_2$ we obtain

$$\begin{aligned} \Phi'(t) &\leq 2\delta\Phi + 4pK_2^2 \int_{\Omega} \operatorname{div} \left[(u^2 + v^2)^{2p-1} (u\nabla u + v\nabla v) \right] dx \\ &\quad - 8p(2p-1)K_2^2 \int_{\Omega} (u^2 + v^2)^{2p-2} (u\nabla u + v\nabla v) (u\nabla u + v\nabla v) dx \\ &\quad - 4pK_2^2 \int_{\Omega} (u^2 + v^2)^{2p-1} (|\nabla u|^2 + |\nabla v|^2) dx + 4pC_0K_2^3 \int_{\Omega} (u^2 + v^2)^{3p} dx. \end{aligned}$$

Now by divergence theorem and boundary conditions in (1.1) and assumption (1.2) we get

$$\begin{aligned} (2.5) \quad \Phi'(t) &\leq 2\delta\Phi - 4pK_2^2\alpha \int_{\partial\Omega} (u^2 + v^2)^{2p} ds \\ &\quad - 8p(2p-1)K_2^2 \int_{\Omega} (u^2 + v^2)^{2p-2} (u\nabla u + v\nabla v) (u\nabla u + v\nabla v) dx \\ &\quad - 4pK_2^2 \int_{\Omega} (u^2 + v^2)^{2p-1} (|\nabla u|^2 + |\nabla v|^2) dx + 4pC_0K_2^3 \int_{\Omega} (u^2 + v^2)^{3p} dx. \end{aligned}$$

Now we neglect the second (negative) term in (2.5), and we use the inequality $(u^2 + v^2)(|\nabla u|^2 + |\nabla v|^2) \geq (u\nabla u + v\nabla v)(u\nabla u + v\nabla v)$ in the fourth term to have

$$\begin{aligned} (2.6) \quad \Phi'(t) &\leq 2\delta\Phi - 4p(4p-1)K_2^2 \int_{\Omega} (u^2 + v^2)^{2p-2} (u\nabla u + v\nabla v) (u\nabla u + v\nabla v) dx \\ &\quad + 4pC_0K_2^3 \int_{\Omega} (u^2 + v^2)^{3p} dx. \end{aligned}$$

For simplicity we define

$$(2.7) \quad V = (u^2 + v^2)^p,$$

then

$$(2.8) \quad V_i V_i = 4p^2(u^2 + v^2)^{2p-2}(u\nabla u + v\nabla v)(u\nabla u + v\nabla v).$$

By plugging (2.7) and (2.8) in (2.6) we have

$$(2.9) \quad \Phi'(t) \leq 2\delta\Phi - \left(4 - \frac{1}{p}\right)K_2^2 \int_{\Omega} V_i V_i + 4pC_0K_2^3 \int_{\Omega} V^3 dx.$$

In the third term in (2.9) we use the inequality

$$\int_{\Omega} V^3 dx \leq \left\{ \frac{3}{2\rho_0} \int_{\Omega} V^2 dx + \left(1 + \frac{d}{\rho_0}\right) \left(\int_{\Omega} V^2 dx\right)^{\frac{1}{2}} \left(\int_{\Omega} |\nabla V|^2 dx\right)^{\frac{1}{2}} \right\}^{\frac{3}{2}}$$

with

$$\rho_0 = \min_{\partial\Omega}(x_i \cdot \nu_i), \quad d^2 = \max_{\Omega}(x_i \cdot x_i),$$

obtained by application of a Sobolev type inequality derived in [8] Lemma A.2, valid only in a convex domain $\subset R^3$ we have

$$(2.10) \quad K_2^3(t) \int_{\Omega} V^3 dx \leq \left\{ \frac{3}{2\rho_0} K_2^2 \int_{\Omega} V^2 dx + \left(1 + \frac{d}{\rho_0}\right) \left(\frac{1}{\tau} K_2^2 \int_{\Omega} V^2 dx\right)^{\frac{1}{2}} \left(\tau K_2^2 \int_{\Omega} |\nabla V|^2 dx\right)^{\frac{1}{2}} \right\}^{\frac{3}{2}}$$

$\tau > 0$ a suitable constant to be chosen later on.

In (2.10) we use the basic inequality $(a + b)^{\frac{3}{2}} \leq \sqrt{2} \left(a^{\frac{3}{2}} + b^{\frac{3}{2}}\right)$ and we obtain

$$(2.11) \quad K_2^3(t) \int_{\Omega} V^3 dx \leq \sqrt{2} \left(\frac{3}{2\rho_0}\right)^{\frac{3}{2}} \Phi^{\frac{3}{2}} + \sqrt{2} \left(1 + \frac{d}{\rho_0}\right)^{\frac{3}{2}} \left(\frac{1}{\tau}\Phi\right)^{\frac{3}{4}} \left(\tau K_2^2 \int_{\Omega} |\nabla V|^2 dx\right)^{\frac{3}{4}}.$$

Now we use the inequality $a^r b^s \leq ra + sb$, $r + s = 1$, we have

$$(2.12) \quad K_2^3 \int_{\Omega} V^3 dx \leq \frac{3^{\frac{3}{2}}}{2\rho_0^{\frac{3}{2}}} \Phi^{\frac{3}{2}} + \frac{1}{2^{\frac{3}{2}}\tau^3} \left(1 + \frac{d}{\rho_0}\right)^{\frac{3}{2}} \Phi^3 + \left(\frac{3}{2^{\frac{3}{2}}}\right) \left(1 + \frac{d}{\rho_0}\right)^{\frac{3}{2}} \tau K_2^2 \int_{\Omega} V_i V_i dx = C_1 \Phi^{\frac{3}{2}} + C_2 \Phi^3 + C_3 \tau K_2^2 \int_{\Omega} V_i V_i dx$$

with

$$(2.13) \quad C_1 = \frac{3^{\frac{3}{2}}}{2\rho_0^{\frac{3}{2}}}, \quad C_2 = \frac{1}{2^{\frac{3}{2}}\tau^3} \left(1 + \frac{d}{\rho_0}\right)^{\frac{3}{2}}, \quad C_3 = \frac{3}{2^{\frac{3}{2}}} \left(1 + \frac{d}{\rho_0}\right)^{\frac{3}{2}}.$$

Inserting (2.12) in (2.9) we obtain

$$(2.14) \quad \Phi'(t) \leq 2\delta\Phi - \left(4 - \frac{1}{p} - 4pC_3C_0\tau\right) K_2^2 \int_{\Omega} V_i V_i + 4pC_1C_0\Phi^{\frac{3}{2}} + 4pC_2C_0\Phi^3.$$

By choosing $\tau = \frac{4 - \frac{1}{p}}{4pC_3C_0}$, we obtain

$$(2.15) \quad \Phi'(t) \leq \xi_1\Phi + \xi_2\Phi^{\frac{3}{2}} + \xi_3\Phi^3 = \phi(\Phi),$$

with $\xi_1 = 2\delta$, $\xi_2 = 4pC_1C_0$, $\xi_3 = 4pC_2C_0$.

Then if Φ blows up at time t^* , there exists a time t_0 such that $\Phi(t) > \Phi(0)$ and by integration from 0 to t^* , we obtain the following lower bound for t^*

$$\int_{\Phi(0)}^{\infty} \frac{d\eta}{\phi(\eta)} \leq t^* - t_0 \leq t^*$$

which is the desired lower bound (2.4).

3. Upper bound. We seek in this section an upper bound for the blow up time t^* by defining the following auxiliary function

$$(3.1) \quad \chi(t) := \chi_1(t) + \chi_2(t) = K_1(t)^{\frac{1}{q-1}} \int_{\Omega} u\varphi_1 dx + K_2(t)^{\frac{1}{q-1}} \int_{\Omega} v\varphi_1 dx,$$

where φ_1 is the first eigenfunction of the following membrane problem

$$(3.2) \quad \begin{cases} \Delta\varphi_1 + \lambda_1\varphi_1 = 0, & \varphi_1 > 0, & \text{in } \Omega, \\ \frac{\partial\varphi_1}{\partial\nu} = -\beta\varphi_1, & & \text{on } \partial\Omega \end{cases}$$

with

$$(3.3) \quad \int_{\Omega} \varphi_1 dx = 1$$

with β in (1.1) satisfying (1.2). We prove the following

Theorem 3.1. *Let (u, v) be the solution of (1.1), in a bounded domain Ω in \mathbb{R}^N . Let $K(t) = \min\{K_1(t), K_2(t)\}$. Assume that*

$$(3.4) \quad f_i(s) \geq s^q \quad q > 1, \quad i = 1, 2,$$

and

$$(3.5) \quad \frac{K'_i}{K_i} \geq \tilde{\gamma},$$

with $\tilde{\gamma}$ a positive constant. Assume moreover that there exists a constant $\bar{\gamma}$ with $0 < \bar{\gamma} \leq 1$ such that

$$(3.6) \quad \frac{K}{K_i} \geq \bar{\gamma}.$$

If (u, v) becomes unbounded in χ measure at some finite time t^* , then t^* is bounded from above by T with

$$(3.7) \quad T = \int_{\chi(0)}^{\infty} \frac{d\eta}{\gamma_2 \eta^q - \gamma_1 \eta},$$

where

$$\gamma_1 = \lambda_1 - \frac{\tilde{\gamma}}{q-1}, \quad \gamma_2 = 2^{1-q} \bar{\gamma}^{\frac{q}{q-1}},$$

provided

$$(3.8) \quad \gamma_2 \chi^{q-1}(0) > \gamma_1.$$

Proof. By derivative of (3.1) we have

$$(3.9) \quad \begin{aligned} \chi'(t) = & \frac{1}{q-1} \left(\frac{K'_1}{K_1} \right) K_1^{\frac{1}{q-1}} \int_{\Omega} u \varphi_1 dx + \frac{1}{q-1} \left(\frac{K'_2}{K_2} \right) K_2^{\frac{1}{q-1}} \int_{\Omega} v \varphi_1 dx \\ & + K_1^{\frac{1}{q-1}} \int_{\Omega} u_t \varphi_1 dx + K_2^{\frac{1}{q-1}} \int_{\Omega} v_t \varphi_1 dx. \end{aligned}$$

By applying in (3.9) the condition (3.5) we obtain

$$(3.10) \quad \begin{aligned} \chi'(t) \geq & \frac{\tilde{\gamma}}{q-1} \chi + K_1^{\frac{1}{q-1}} \int_{\Omega} \Delta u \varphi_1 dx + K_2^{\frac{1}{q-1}} \int_{\Omega} \Delta v \varphi_1 dx \\ & + K_1^{\frac{q}{q-1}} \int_{\Omega} f_1 \varphi_1 dx + K_2^{\frac{q}{q-1}} \int_{\Omega} f_2 \varphi_1 dx, \end{aligned}$$

We compute

$$\int_{\Omega} \Delta u \varphi_1 dx = \int_{\Omega} \operatorname{div}[\nabla(u\varphi_1)] dx - \int_{\Omega} u \Delta \varphi_1 dx - 2 \int_{\Omega} \nabla u \nabla \varphi_1 dx.$$

By the equation in (3.2), the divergence theorem and boundary condition in (3.2) we obtain

$$(3.11) \quad \int_{\Omega} \Delta u \varphi_1 dx = (\beta - \alpha) \int_{\partial\Omega} u \varphi_1 ds - \lambda_1 \int_{\Omega} u \varphi_1 dx \geq -\lambda_1 \int_{\Omega} u \varphi_1 dx$$

analogously, we obtain

$$(3.12) \quad \int_{\Omega} \Delta v \varphi_1 dx = -\lambda_1 \int_{\Omega} v \varphi_1 dx.$$

By inserting (3.11) and (3.12) in (3.10) we have

$$(3.13) \quad \begin{aligned} \chi'(t) &\geq \frac{\tilde{\gamma}}{q-1} \chi - \lambda_1 \left[K_1^{\frac{1}{q-1}} \int_{\Omega} u \varphi_1 dx + K_2^{\frac{1}{q-1}} \int_{\Omega} v \varphi_1 dx \right] \\ &\quad + K_1^{\frac{q}{q-1}} \int_{\Omega} f_1 \varphi_1 dx + K_2^{\frac{q}{q-1}} \int_{\Omega} f_2 \varphi_1 dx \\ &= - \left(\lambda_1 - \frac{\tilde{\gamma}}{q-1} \right) \chi + K_1^{\frac{q}{q-1}} \int_{\Omega} u^q \varphi_1 dx + K_2^{\frac{q}{q-1}} \int_{\Omega} v^q \varphi_1 dx. \end{aligned}$$

In the last two terms of (3.13) we used (3.4), and by expression of $K(t)$ we obtain

$$(3.14) \quad \chi'(t) \geq -\gamma_1 \chi + K^{\frac{q}{q-1}} \int_{\Omega} (u^q + v^q) \varphi_1 dx,$$

with $\gamma_1 = \lambda_1 - \frac{\tilde{\gamma}}{q-1}$.

To estimate $\int_{\Omega} (u^q + v^q) \varphi_1 dx$ we use Hölder inequality and (3.3)

$$(3.15) \quad \begin{cases} \int_{\Omega} u^q \varphi_1 dx \geq \left(\int_{\Omega} u \varphi_1 dx \right)^q \\ \int_{\Omega} v^q \varphi_1 dx \geq \left(\int_{\Omega} v \varphi_1 dx \right)^q \end{cases}$$

By substitution of (3.15) in (3.14) we find

$$(3.16) \quad \chi'(t) \geq -\gamma_1 \chi + K^{\frac{q}{q-1}} \left\{ \left(\int_{\Omega} u \varphi_1 dx \right)^q + \left(\int_{\Omega} v \varphi_1 dx \right)^q \right\}$$

$$= -\gamma_1\chi + \left(\frac{K}{K_1}\right)^{\frac{q}{q-1}} \chi_1^q + \left(\frac{K}{K_2}\right)^{\frac{q}{q-1}} \chi_2^q.$$

By inserting in the last term of (3.16) the assumption (3.6), we obtain

$$(3.17) \quad \chi'(t) \geq -\gamma_1\chi + \bar{\gamma}^{\frac{q}{q-1}}(\chi_1^q + \chi_2^q).$$

Using in (3.17) the inequality $\chi_1^q + \chi_2^q \geq 2^{1-q}(\chi_1 + \chi_2)^q$ we obtain

$$(3.18) \quad \chi'(t) \geq -\gamma_1\chi + \gamma_2\chi^q.$$

with $\gamma_2 = 2^{1-q}\bar{\gamma}^{\frac{q}{q-1}}$.

Since (3.8) holds we conclude that $\chi(t)$ is an increasing function of $t \in (0, t^*)$ and that by integration of (3.18) from 0 to t we find

$$(3.19) \quad \int_{\chi(0)}^{\chi(t)} \frac{d\eta}{\gamma_2\eta^q - \gamma_1\eta} \geq t.$$

and we obtain the upper bound for t^* :

$$t^* \leq \int_{\chi(0)}^{\infty} \frac{d\eta}{\eta(\gamma_2\eta^{q-1} - \gamma_1)}.$$

Acknowledgments. The author thanks G.Philippin and S.Vernier Piro for useful discussions on the subject.

REFERENCES

- [1] J. M. BALL. Remarks on blow-up and nonexistence theorems for nonlinear evolution equations. *Quart. J. Math. Oxford* **28** (1977), 473–486.
- [2] H. KIELHÖFER. Halbgruppen und semilineare Anfangs-randwert-probleme. *Manuscripta Math.* **12** (1974), 121–152.
- [3] M. MARRAS. Bounds for blow-up time in nonlinear parabolic systems under various boundary conditions. *Numer. Funct. Anal. Optim.* **32**, (2011), 453–468.

- [4] M. MARRAS, S. VERNIER PIRO. Blow-up phenomena in reaction-diffusion systems. In: Special Issue of Discrete and Continuous Dynamical Systems on Nonlinear Elliptic and Parabolic Problems, 2011 (to appear).
- [5] L. E. PAYNE, G. A. PHILIPPIN, P. W. SCHAEFER. Bounds for blow-up time in nonlinear parabolic problems. *J. Math. Anal. Appl.* **338** (2008), 438–447.
- [6] L. E. PAYNE, G. A. PHILIPPIN, P. W. SCHAEFER. Blow-up phenomena for some nonlinear parabolic problems. *Nonlinear Anal.* **69** (2008), 3495–3502.
- [7] L. E. PAYNE, G. A. PHILIPPIN, S. VERNIER-PIRO. Blow-up phenomena for a semilinear heat equation with nonlinear boundary condition I. *Z. Angew. Math. Phys.* **61** (2010), 971–978.
- [8] L. E. PAYNE, G. A. PHILIPPIN, S. VERNIER-PIRO. Blow-up phenomena for a semilinear heat equation with nonlinear boundary condition II. *Nonlinear Anal.* **73** (2010), 971–978.
- [9] L. E. PAYNE, P. W. SCHAEFER. Blow-up phenomena for some nonlinear parabolic systems. *Int. J. Pure Appl. Math.* **42** (2008), 193–202.
- [10] P. QUITTNER, P. SOUPLET. Superlinear Parabolic Problems. Blow-up, Global Existence and Steady States, Birkhäuser Advanced Texts, Basel, 2007.
- [11] B. STRAUGHAN. Explosive Instabilities in Mechanics. Springer, Berlin, 1998.
- [12] J. L. VAZQUEZ. The problems of blow-up for nonlinear heat equations. Complete blow-up and avalanche formation. *Rend. Mat. Acc. Lincei ser. 9* **15** (2004), 281–300.
- [13] F. B. WEISSLER. Local existence and nonexistence for semilinear parabolic equations in L^p . *Indiana Univ. Math. J.* **29** (1980), 79–102.
- [14] F. B. WEISSLER. Existence and nonexistence of global solutions for a heat equation. *Israel J. Math.* **38**, 1–2 (1981), 29–40.

Dipartimento di Matematica e Informatica
Universita' di Cagliari
Italy
e-mail: mmarras@unica.it

Received July 4, 2011