

Provided for non-commercial research and educational use.  
Not for reproduction, distribution or commercial use.

# Serdica

## Mathematical Journal

### Сердика

### Математическо списание

---

The attached copy is furnished for non-commercial research and education use only.  
Authors are permitted to post this version of the article to their personal websites or institutional repositories and to share with other researchers in the form of electronic reprints.  
Other uses, including reproduction and distribution, or selling or licensing copies, or posting to third party websites are prohibited.

For further information on  
Serdica Mathematical Journal  
which is the new series of  
Serdica Bulgaricae Mathematicae Publicationes  
visit the website of the journal <http://www.math.bas.bg/~serdica>  
or contact: Editorial Office  
Serdica Mathematical Journal  
Institute of Mathematics and Informatics  
Bulgarian Academy of Sciences  
Telephone: (+359-2)9792818, FAX:(+359-2)971-36-49  
e-mail: [serdica@math.bas.bg](mailto:serdica@math.bas.bg)

## RELIABILITY FOR BETA MODELS

Saralees Nadarajah

*Communicated by N. M. Yanev*

ABSTRACT. In the area of stress-strength models there has been a large amount of work as regards estimation of the reliability  $R = \Pr(X_2 < X_1)$  when  $X_1$  and  $X_2$  are independent random variables belonging to the same univariate family of distributions. The algebraic form for  $R = \Pr(X_2 < X_1)$  has been worked out for the majority of the well-known distributions including Normal, uniform, exponential, gamma, weibull and pareto. However, there are still many other distributions for which the form of  $R$  is not known. We have identified at least some 30 distributions with no known form for  $R$ . In this paper we consider some of these distributions and derive the corresponding forms for the reliability  $R$ . The calculations involve the use of various special functions.

**1. Introduction.** In the context of reliability the stress–strength model describes the life of a component which has a random strength  $X_1$  and is subjected to random stress  $X_2$ . The component fails at the instant that the stress applied to

---

2000 *Mathematics Subject Classification*: 33C90, 62E99.

*Key words*: Beta distributions, Hypergeometric functions, Incomplete beta function, Reliability.

it exceeds the strength and the component will function satisfactorily whenever  $X_1 > X_2$ . Thus  $R = \Pr(X_2 < X_1)$  is a measure of component reliability. It has applications in many areas. For example, if  $X_2$  represents the maximum chamber pressure generated by ignition of a solid propellant and  $X_1$  represents the strength of the rocket chamber then  $R$  is the probability of successful firing of the engine. Another example is when  $X_2$  represents the diameter of a shaft and  $X_1$  represents the diameter of a bearing that is to be mounted on the shaft – here  $R$  is the probability that the bearing fits without interference. Because of these applications, the calculation and the estimation of  $R = \Pr(X_2 < X_1)$  is important. This has been investigated extensively in the literature when  $X_1$  and  $X_2$  are independent random variables belonging to the same univariate family of distributions. The algebraic form for  $R$  has been worked out for the majority of the well-known distributions in their standard forms. These include Normal, uniform, exponential, gamma, weibull and the pareto distributions. However, we have identified many other distributions including extensions of the above distributions for which the form of  $R$  is not known. In this paper we consider the class of beta distributions and derive the corresponding forms for  $R$ .

We shall assume throughout this paper that  $X_1$  and  $X_2$  are continuous and independent random variables. Let  $f_i$  and  $F_i$  denote, respectively, the probability density function (pdf) and the cumulative distribution function (cdf) of  $X_i$ . With this notation, we can write

$$(1) \quad \begin{aligned} R &= \Pr(X_2 < X_1) \\ &= \int_{-\infty}^{\infty} F_2(x) f_1(x) dx. \end{aligned}$$

The calculations of (1) will make use of the following special functions: the gamma function defined by

$$\Gamma(a) = \int_0^{\infty} z^{a-1} \exp(-z) dz;$$

the beta function defined by

$$B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)};$$

the incomplete beta function ratio defined by

$$(2) \quad I_x(a, b) = \frac{1}{B(a, b)} \int_0^x w^{a-1} (1-w)^{b-1} dw;$$

and, the generalized hypergeometric function defined by

$$(3) \quad {}_pF_q(\alpha_1, \alpha_2, \dots, \alpha_p; \beta_1, \beta_2, \dots, \beta_q; x) = \sum_{k=0}^{\infty} \frac{(\alpha_1)_k (\alpha_2)_k \cdots (\alpha_p)_k x^k}{(\beta_1)_k (\beta_2)_k \cdots (\beta_q)_k k!}.$$

When  $p = 1$  and  $q = 1$ , (3) is known as the confluent hypergeometric function. When  $p = 2$  and  $q = 1$ , (3) is known as the Gauss hypergeometric function. Some properties of the incomplete beta function ratio that we shall need are:

$$(4) \quad I_x(a, b) = \frac{x^a}{aB(a, b)} {}_2F_1(a, 1 - b; 1 + a; x),$$

$$(5) \quad I_x(a, n - a + 1) = \sum_{k=a}^n \binom{n}{k} x^k (1 - x)^{n-k},$$

if  $a$  is an integer,

$$(6) \quad I_x(a, b) = 1 - \sum_{k=1}^a \frac{\Gamma(b + k - 1)}{\Gamma(b)\Gamma(k)} x^{k-1} (1 - x)^b,$$

if  $a$  is an integer,

$$(7) \quad I_x(a, b) = \sum_{k=1}^b \frac{\Gamma(a + k - 1)}{\Gamma(a)\Gamma(k)} x^a (1 - x)^{k-1},$$

if  $b$  is an integer,

and

$$(8) \quad I_x\left(k - \frac{1}{2}, j - \frac{1}{2}\right) = \frac{2}{\pi} \arctan \sqrt{\frac{x}{1-x}} + \sum_{l=1}^{j-1} c_l - \sqrt{x(1-x)} \sum_{l=1}^{k-1} d_l,$$

where

$$c_l = \frac{\Gamma(k + l - 1)}{\Gamma(k - 1/2)\Gamma(l + 1/2)} x^{k-1/2} (1 - x)^{l-1/2},$$

$$d_l = \frac{\Gamma(l)}{\Gamma(l + 1/2)\Gamma(1/2)} x^{l-1}.$$

Some properties of the generalized hypergeometric function that we shall need are:

$$(9) \quad {}_2F_1(a, b; c; 1) = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)},$$

$$c > a + b,$$

$$(10) \quad {}_3F_2(a, b, c; a+1, b+1; 1) = \frac{ab}{a-b}\Gamma(1-c) \left\{ \frac{\Gamma(b)}{\Gamma(1+b-c)} - \frac{\Gamma(a)}{\Gamma(1+a-c)} \right\},$$

$$a \neq b, \quad c \neq 1, \quad c < 2,$$

$$(11) \quad \int_0^1 z^{a-1}(1-z)^{b-1} {}_2F_1(c, d; e; z) dz = B(a, b) {}_3F_2(c, d, a; e, a+b; 1),$$

$$a > 0, \quad b > 0, \quad e + b > c + d$$

and

$$(12) \quad \int_0^x z^{a-1}(x-z)^{b-1} {}_2F_1(c, d; e; z) dz = B(a, b)x^{a+b-1} {}_3F_2(c, d, a; e, a+b; x),$$

$$x > 0, \quad a > 0, \quad b > 0.$$

We shall also need the following properties:

$$(13) \quad \int_0^\infty \frac{\arctan z}{z} \left( \frac{z^p}{1+z^{2p}} \right)^{2q} dz = \frac{\pi^{3/2}}{2^{2q+2p}} \frac{\Gamma(q)}{\Gamma(q+1/2)},$$

$$q > 0,$$

$$(14) \quad \int_0^x z^{d-1}(z+a)^c dz = \frac{a^c x^d}{d} {}_2F_1\left(-c, d; 1+d; -\frac{x}{a}\right),$$

$$d > 0$$

and

$$(15) \quad \int_0^x z^{b-1}(x-z)^{a-1} \exp(cz) dz = B(a, b)x^{a+b-1} {}_1F_1(b; a+b; cx),$$

$$a > 0, \quad b > 0.$$

Further properties of the special functions being used can be found in Prudnikov et al. [9] and Gradshteyn and Ryzhik [3].

**2. Standard Beta Distribution**

For the standard form of the beta distribution, the pdf and the cdf of  $X_i$  are

$$(16) \quad f_i(x) = \frac{1}{B(a_i, b_i)} x^{a_i-1} (1-x)^{b_i-1}$$

and

$$(17) \quad F_i(x) = I_x(a_i, b_i),$$

(where  $0 \leq x \leq 1$ ), respectively. These distributions are very versatile and a variety of uncertainties can be usefully modeled by them. In recent years beta distributions have attracted applications in activity in PERT analysis, breakage models, communication theory, gas absorption, hydrology, particle size distributions, photovoltaic system analysis, sea-state reflectivity, solar radiation and other areas. For (16) and (17), the reliability given by (1) takes the form:

$$(18) \quad R = \int_0^1 \frac{I_x(a_2, b_2)}{B(a_1, b_1)} x^{a_1-1} (1-x)^{b_1-1} dx.$$

This can be evaluated by applying (4) to re-express  $I_x(a_2, b_2)$  and then using (11) to calculate the integral:

$$(19) \quad R = \frac{1}{a_2 B(a_1, b_1) B(a_2, b_2)} \int_0^1 x^{a_1+a_2-1} (1-x)^{b_1-1} {}_2F_1(a_2, 1-b_2; 1+a_2; x) dx$$

$$= \frac{B(a_1 + a_2, b_1) {}_3F_2(a_2, 1 - b_2, a_1 + a_2; 1 + a_2, a_1 + a_2 + b_1; 1)}{a_2 B(a_1, b_1) B(a_2, b_2)}.$$

This expression for  $R$  can be reduced to elementary forms for the particular choices of the parameters  $a_i$  and  $b_i$  considered below.

Case 1: If  $a_1 + a_2 + b_1 + b_2 = 1$  then, using (9), we have

$$\begin{aligned} & {}_3F_2(a_2, 1 - b_2, a_1 + a_2; 1 + a_2, a_1 + a_2 + b_1; 1) \\ &= {}_2F_1(a_2, a_1 + a_2; 1 + a_2; 1) \\ &= \frac{\Gamma(1 + a_2) \Gamma(1 - a_1 - a_2)}{\Gamma(1 - a_1)} \\ &= \frac{1 - a_1}{B(1 + a_2, 1 - a_1 - a_2)}; \end{aligned}$$

so, (19) reduces to:

$$R = \frac{(1 - a_1) B(a_1 + a_2, b_1)}{a_2 B(a_1, b_1) B(a_2, b_2) B(1 + a_2, 1 - a_1 - a_2)}.$$

Case 2: If  $a_1 = 1$  then, again using (9), we have

$$\begin{aligned} & {}_3F_2(a_2, 1 - b_2, a_1 + a_2; 1 + a_2, a_1 + a_2 + b_1; 1) \\ &= {}_2F_1(a_2, 1 - b_2; 1 + a_2 + b_1; 1) \\ &= \frac{\Gamma(1 + a_2 + b_1) \Gamma(b_1 + b_2)}{\Gamma(1 + b_1) \Gamma(a_2 + b_1 + b_2)} \\ &= \frac{B(1 + a_2 + b_1, b_1 + b_2)}{B(1 + b_1, a_2 + b_1 + b_2)}; \end{aligned}$$

so, (19) reduces to:

$$R = \frac{B(a_1 + a_2, b_1) B(1 + a_2 + b_1, b_1 + b_2)}{a_2 B(a_1, b_1) B(a_2, b_2) B(1 + b_1, a_2 + b_1 + b_2)}.$$

Case 3: If  $b_1 = 1$  then, using (10), we have

$$\begin{aligned} & {}_3F_2(a_2, 1 - b_2, a_1 + a_2; 1 + a_2, a_1 + a_2 + b_1; 1) \\ &= \frac{a_2(a_1 + a_2)}{(-a_1)} \Gamma(b_2) \left\{ \frac{\Gamma(a_1 + a_2)}{\Gamma(a_1 + a_2 + b_2)} - \frac{\Gamma(a_2)}{\Gamma(a_2 + b_2)} \right\} \\ &= \frac{a_2(a_1 + a_2)}{a_1} \{B(a_2, b_2) - B(a_1 + a_2, b_2)\}; \end{aligned}$$

so, (19) reduces to:

$$R = \frac{(a_1 + a_2) B(a_1 + a_2, b_1) \{B(a_2, b_2) - B(a_1 + a_2, b_2)\}}{a_1 B(a_1, b_1) B(a_2, b_2)}.$$

Case 4: If  $a_2 = a$  and  $b_2 = n - a + 1$  (where  $a$  is an integer) then, substituting (5) into (18), we have

$$\begin{aligned} R &= \sum_{k=a}^n \frac{1}{B(a_1, b_1)} \binom{n}{k} \int_0^1 x^{a_1+k-1} (1-x)^{b_1+n-k-1} dx \\ &= \sum_{k=a}^n \binom{n}{k} \frac{B(a_1 + k, b_1 + n - k)}{B(a_1, b_1)}. \end{aligned}$$

Case 5: If  $a_2$  is an integer then, substituting (6) into (18), we have

$$\begin{aligned}
 R &= 1 - \sum_{k=1}^{a_2} \frac{1}{B(a_1, b_1)} \frac{\Gamma(b_2 + k - 1)}{\Gamma(b_2)\Gamma(k)} \int_0^1 x^{a_1+k-2}(1-x)^{b_1+b_2-1} dx \\
 &= 1 - \sum_{k=1}^{a_2} \frac{B(a_1 + k - 1, b_1 + b_2)}{(b_2 + k - 1) B(a_1, b_1) B(b_2, k)}.
 \end{aligned}$$

In the particular case  $a_2 = 1$ ,

$$R = 1 - \frac{B(a_1, b_1 + b_2)}{B(a_1, b_1)}.$$

Case 6: Similarly, if  $b_2$  is an integer, substituting (7) into (18), we have

$$R = \sum_{k=1}^{b_2} \frac{B(a_1 + a_2, b_1 + k - 1)}{(a_2 + k - 1) B(a_1, b_1) B(a_2, k)}.$$

In the particular case  $b_2 = 1$ ,

$$R = \frac{B(a_1 + a_2, b_1)}{B(a_1, b_1)}.$$

Case 7: Finally, if  $a_2 = k - 1/2$  and  $b_2 = j - 1/2$ , substituting (8) into (18), we have

$$\begin{aligned}
 R &= \frac{2}{\pi B(a_1, b_1)} \int_0^1 \arctan \sqrt{\frac{x}{1-x}} x^{a_1-1} (1-x)^{b_1-1} dx \\
 &\quad + \sum_{l=1}^{j-1} \frac{B(a_1 + k - 1/2, b_1 + l - 1/2)}{(k + l - 1) B(k - 1/2, l + 1/2) B(a_1, b_1)} \\
 (20) \quad &\quad - \sum_{l=1}^{k-1} \frac{B(a_1 + l - 1/2, b_1 + 1/2)}{l B(l + 1/2, 1/2) B(a_1, b_1)}.
 \end{aligned}$$

On substituting  $y = \sqrt{x/(1-x)}$ , the integral term reduces to

$$(21) \quad 2 \int_0^\infty \frac{y^{2a_1-1} \arctan y}{(1+y^2)^{a_1+b_1}} dy.$$



This integral cannot be simplified further for general  $a_1$  and  $b_1$ . However, in the particular case  $a_1 = b_1 = c$  (say), using (13) with  $p = 1$  and  $q = c$ , (21) can be reduced to

$$\frac{\pi^{3/2}}{2^{2c+1}} \frac{\Gamma(c)}{\Gamma(c + 1/2)}.$$

Substituting this into (20), we get

$$R = \frac{\sqrt{\pi}\Gamma(2c)}{4^c\Gamma(c)\Gamma(c + 1/2)} + \sum_{l=1}^{j-1} \frac{B(c + k - 1/2, c + l - 1/2)}{(k + l - 1)B(k - 1/2, l + 1/2)B(c, c)} - \sum_{l=1}^{k-1} \frac{B(c + l - 1/2, c + 1/2)}{lB(l + 1/2, 1/2)B(c, c)}$$

for the particular case  $a_1 = b_1 = c$ .

**3. Uniform distribution.** If  $X_i$  have the uniform distribution then the pdf and the cdf are

$$(22) \quad f_i(x) = \frac{1}{d_i - c_i}$$

and

$$(23) \quad F_i(x) = \frac{x - c_i}{d_i - c_i},$$

(where  $c_i \leq x \leq d_i$ ), respectively. These distributions have found extensive applications in life testing and traffic flow modeling. The standard forms of (22)–(23) when  $c_i = 0$  and  $d_i = 1$  are particular cases of the beta distribution in (16)–(17). The form of the reliability for (22)–(23) is easy to calculate as shown below:

$$R = \int_{\max(c_1, c_2)}^{\min(d_1, d_2)} \frac{x - c_2}{d_2 - c_2} \frac{1}{d_1 - c_1} dx + \int_{\min(d_1, d_2)}^{d_1} \frac{1}{d_1 - c_1} dx$$

$$= \frac{\{\min(d_1, d_2)\}^2 - \{\max(c_1, c_2)\}^2 + 2c_2 \{\max(c_1, c_2) - \min(d_1, d_2)\}}{2(d_1 - c_1)(d_2 - c_2)} + \frac{d_1 - \min(d_1, d_2)}{d_1 - c_1}.$$

In the standard case  $c_1 = c_2 = 0$  and  $d_1 = d_2 = 1$ , we have  $R = 1/2$  as expected (this is also true if  $c_1 = c_2$  and  $d_1 = d_2$ ).

**4. Power Function Distribution.** For a power function distribution the pdf and the cdf of  $X_i$  are

$$(24) \quad f_i(x) = a_i \left( \frac{x - c_i}{d_i - c_i} \right)^{a_i - 1}$$

and

$$(25) \quad F_i(x) = (d_i - c_i) \left( \frac{x - c_i}{d_i - c_i} \right)^{a_i},$$

(where  $c_i \leq x \leq d_i$ ), respectively. These distributions have found recent applications in electrical component reliability (Meniconi and Barry, [6]) and mineral resource prediction (Shen and Zhao, [11]). The standard forms of (24)–(25) when  $c_i = 0$  and  $d_i = 1$  are particular cases of the beta distribution in (16)–(17). The reliability for (24)–(25) takes the form:

$$(26) \quad R = a_1 \int_{\max(c_1, c_2)}^{\min(d_1, d_2)} (d_2 - c_2) \left( \frac{x - c_2}{d_2 - c_2} \right)^{a_2} \left( \frac{x - c_1}{d_1 - c_1} \right)^{a_1 - 1} dx + a_1 \int_{\min(d_1, d_2)}^{d_1} \left( \frac{x - c_1}{d_1 - c_1} \right)^{a_1 - 1} dx.$$

The second term on the right of (26) is:

$$1 - F_1(\min(d_1, d_2)) = 1 - (d_1 - c_1) \left( \frac{\min(d_1, d_2) - c_1}{d_1 - c_1} \right)^{a_1}.$$

The first term on the right of (26), say  $J$ , can be evaluated using (14). If  $c_1 > c_2$  then, substituting  $y = x - c_1$ , we see

$$J = \frac{a_1}{(d_1 - c_1)^{a_1 - 1} (d_2 - c_2)^{a_2 - 1}} \int_0^{\min(d_1, d_2) - c_1} (y + c_1 - c_2)^{a_2} y^{a_1 - 1} dy$$

which reduces to

$$J = \frac{(c_1 - c_2)^{a_2} \{\min(d_1, d_2) - c_1\}^{a_1}}{(d_1 - c_1)^{a_1 - 1} (d_2 - c_2)^{a_2 - 1}} {}_2F_1 \left( -a_2, a_1; 1 + a_1; \frac{\min(d_1, d_2) - c_1}{c_2 - c_1} \right)$$

on applying (14). Similarly, if  $c_2 > c_1$  then

$$J = \frac{a_1(c_2 - c_1)^{a_1 - 1} \{\min(d_1, d_2) - c_2\}^{1 + a_2}}{(1 + a_2)(d_1 - c_1)^{a_1 - 1} (d_2 - c_2)^{a_2 - 1}} {}_2F_1 \left( 1 - a_1, 1 + a_2; 2 + a_2; \frac{\min(d_1, d_2) - c_2}{c_1 - c_2} \right).$$

If  $c_1 = c_2 = c$  (say) then it is easy to see

$$J = \frac{a_1 \{\min(d_1, d_2) - c\}^{a_1 + a_2}}{(a_1 + a_2) (d_1 - c)^{a_1 - 1} (d_2 - c)^{a_2 - 1}}.$$

**5. Arc-sine distribution.** For an arc-sine distribution the pdf and the cdf of  $X_i$  are

$$(27) \quad f_i(x) = \frac{1}{\pi \sqrt{(4/c_i^2) - x^2}}$$

and

$$(28) \quad F_i(x) = \frac{1}{2} + \frac{1}{\pi} \arcsin \left( \frac{|c_i| x}{2} \right),$$

(where  $-2/|c_i| \leq x \leq 2/|c_i|$ ), respectively. The arc-sine distribution arises naturally in statistical communication theory; see Lee [5, Chapter 6] and Middleton [7, Chapter 14], where (27) is used as a model for the amplitude of a periodic signal in thermal noise and the limiting spectral density function of a high-index-angle modulated carrier, respectively. The arc-sine distribution arises also in the study of the simple random walk. The standard forms of (27) and (28) when  $c_i = 2$  are particular cases of the beta distribution in (16)–(17). The reliability, (1), for (27)–(28) can be written as:

$$(29) \quad R = \frac{1}{2\pi} \int_{-\delta}^{\delta} \left( \frac{4}{c_1^2} - x^2 \right)^{-1/2} dx + \frac{1}{\pi^2} \int_{-\delta}^{\delta} \arcsin \left( \frac{|c_2| x}{2} \right) \left( \frac{4}{c_1^2} - x^2 \right)^{-1/2} dx + \frac{1}{\pi} \int_{\delta}^{2/|c_1|} \left( \frac{4}{c_1^2} - x^2 \right)^{-1/2} dx,$$

where  $\delta = 2 \min(1/|c_1|, 1/|c_2|)$ . The first and the third terms on the right of (29) are equal to:

$$\frac{1}{2} \{F_1(\delta) - F_1(-\delta)\} = \frac{1}{\pi} \arcsin \left[ \frac{|c_1|}{\max(|c_1|, |c_2|)} \right]$$

and

$$1 - F_1(\delta) = \frac{1}{2} - \frac{1}{\pi} \arcsin \left[ \frac{|c_1|}{\max(|c_1|, |c_2|)} \right],$$

respectively. The second term on the right of (29) is zero because its integrand is an odd function. Thus for the arc-sine distribution  $R = 1/2$  irrespective of what  $c_1$  and  $c_2$  are.

**6. Generalized Beta distribution.** A four-parameter generalization of the standard beta distribution in (16) and (17) is given by

$$(30) \quad f_i(x) = \frac{1}{(d_i - c_i) B(a_i, b_i)} \left( \frac{x - c_i}{d_i - c_i} \right)^{a_i-1} \left( 1 - \frac{x - c_i}{d_i - c_i} \right)^{b_i-1}$$

and

$$(31) \quad F_i(x) = I_{\frac{x-c_i}{d_i-c_i}}(a_i, b_i),$$

where  $c_i \leq x \leq d_i$ . These contain the uniform and the power function distributions (discussed above) as special cases. Setting  $Y_i = (X_i - c_i)/(d_i - c_i)$ , one can see (30) and (31) reduce to the standard forms. So, if  $c_1 = c_2$  and  $d_1 = d_2$  then  $R = \Pr(X_2 < X_1) = \Pr(Y_2 < Y_1)$  takes the expression given in (19). In the general case,  $R$  can be written as

$$(32) \quad R = \frac{1}{(d_1 - c_1)B(a_1, b_1)} \int_{\max(c_1, c_2)}^{\min(d_1, d_2)} I_{\frac{x-c_2}{d_2-c_2}}(a_2, b_2) \left( \frac{x-c_1}{d_1-c_1} \right)^{a_1-1} \left( 1 - \frac{x-c_1}{d_1-c_1} \right)^{b_1-1} dx$$

$$+ \frac{1}{(d_1 - c_1)B(a_1, b_1)} \int_{\min(d_1, d_2)}^{d_1} \left( \frac{x - c_1}{d_1 - c_1} \right)^{a_1-1} \left( 1 - \frac{x - c_1}{d_1 - c_1} \right)^{b_1-1} dx.$$

The second term on the right of (32) is

$$1 - F_1(\min(d_1, d_2)) = 1 - I_{\frac{\min(d_1, d_2) - c_1}{d_1 - c_1}}(a_1, b_1).$$

On using (4) and substituting  $y = (x - a_2)/(b_2 - a_2)$ , the first term on the right of (32) can be re-written as

$$\frac{(c_2 - c_1)^{a_1-1} (d_1 - c_2)^{b_1-1} (d_2 - c_2)}{a_2 (d_1 - c_1)^{a_1+b_1-1} B(a_1, b_1) B(a_2, b_2)} J,$$

where

$$(33) \quad J = \int_{\delta_1}^{\delta_2} y^{a_2} \left(1 + \frac{d_2 - c_2}{c_2 - c_1} y\right)^{a_1 - 1} \left(1 - \frac{d_2 - c_2}{d_1 - c_2} y\right)^{b_1 - 1} {}_2F_1(a_2, 1 - b_2; 1 + a_2; y) dy$$

with  $\delta_1 = \{\max(c_1, c_2) - c_2\}/(d_2 - c_2)$  and  $\delta_2 = \{\min(d_1, d_2) - c_2\}/(d_2 - c_2)$ . This integral cannot be simplified further for general  $a_1$  and  $b_1$ . However, if we assume that  $a_1 > 1$  and  $b_1 > 1$  are both integers then using binomial expansion we can re-write (33) as

$$J = \sum_{k=0}^{a_1 - 1} \sum_{l=0}^{b_1 - 1} (-1)^l \binom{a_1 - 1}{k} \binom{b_1 - 1}{l} \left(\frac{d_2 - c_2}{c_2 - c_1}\right)^k \left(\frac{d_2 - c_2}{d_1 - c_2}\right)^l J(k, l),$$

where  $J(k, l)$  is the simpler integral

$$(34) \quad J(k, l) = \int_{\delta_1}^{\delta_2} y^{k+l+a_2} {}_2F_1(a_2, 1 - b_2; 1 + a_2; y) dy.$$

On applying (12), (34) can be evaluated as

$$J(k, l) = \frac{\delta_2^{1+k+l+a_2}}{1+k+l+a_2} {}_3F_2(a_2, 1 - b_2, 1 + k + l + a_2; 1 + a_2, 2 + k + l + a_2; \delta_2) - \frac{\delta_1^{1+k+l+a_2}}{1+k+l+a_2} {}_3F_2(a_2, 1 - b_2, 1 + k + l + a_2; 1 + a_2, 2 + k + l + a_2; \delta_1).$$

**7. Non-central Beta distribution.** There are three types of non-central-beta distributions. The one that is most commonly known and studied is the Type I noncentral-beta distribution. For this, the pdf and the cdf of  $X_i$  are

$$(35) \quad f_i(x) = \sum_{k=0}^{\infty} \frac{(\lambda_i/2)^k \exp(-\lambda_i/2) x^{a_i+k-1} (1-x)^{b_i-1}}{k! B(a_i+k, b_i)}$$

and

$$(36) \quad F_i(x) = \sum_{k=0}^{\infty} \frac{(\lambda_i/2)^k \exp(-\lambda_i/2)}{k!} I_x(a_i+k, b_i),$$

(where  $0 \leq x \leq 1$ ), respectively. These distributions have been recently utilized as models in geophysics (Kimball and Scheibner, [4]) and psychophysics (Rousseau and Ennis, [10]). For (35) and (36), the reliability in (1) can be written as:

$$(37) \quad R = \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \frac{\lambda_1^k \lambda_2^l}{2^{k+l} k! l!} \exp\left(-\frac{\lambda_1 + \lambda_2}{2}\right) \int_0^1 \frac{I_x(l+a_2, b_2)}{B(k+a_1, b_1)} x^{k+a_1-1} (1-x)^{b_1-1} dx.$$

The integral on the right has the form of the reliability expression (18) for the standard beta distribution. Thus, using (19), (37) can be reduced to:

$$R = \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \frac{\lambda_1^k \lambda_2^l B(k+l+a_1+a_2, b_1)}{2^{k+l} k! l! (l+a_2) B(k+a_1, b_1) B(l+a_2, b_2)} \exp\left(-\frac{\lambda_1 + \lambda_2}{2}\right) \times {}_3F_2(l+a_2, 1-b_2, k+l+a_1+a_2; 1+l+a_2, k+l+a_1+a_2+b_1; 1).$$

Clearly this is a very complicated expression. A simpler expression can be found if we assume that the  $b_i$  in (35) and (36) are integers. In this case, (35) and (36) reduce to

$$f_i(x) = \exp\{-\lambda_i(1-x)\} \sum_{k=0}^{b_i-1} \sum_{l=1}^{b_i-k} \frac{\lambda_i^k}{k!} \binom{a_i+b_i-l-1}{b_i-k-l} x^{a_i+k-1} (1-x)^{b_i-l-1} \times \{-\lambda_i x^2 + (l-k+\lambda_i-a_i-b_i)x+k+a_i\}$$

and

$$F_i(x) = \exp\{-\lambda_i(1-x)\} \sum_{k=0}^{b_i-1} \sum_{l=1}^{b_i-k} \frac{\lambda_i^k}{k!} \binom{a_i+b_i-l-1}{b_i-k-l} x^{a_i+k} (1-x)^{b_i-l},$$

respectively (see Nicholson [8]). Thus the corresponding form for the reliability can be written as

$$(38) \quad R = \exp\{-(\lambda_1 + \lambda_2)\} \sum_{k=0}^{b_2-1} \sum_{l=1}^{b_2-k} \sum_{m=0}^{b_1-1} \sum_{n=1}^{b_1-m} \frac{\lambda_1^m \lambda_2^k}{m! k!} \binom{a_2+b_2-l-1}{b_2-k-l} \binom{a_1+b_1-n-1}{b_1-m-n} \times \{-\lambda_1 I_1 + (n-m+\lambda_1-a_1-b_1) I_2 + (m+a_1) I_3\},$$

where

$$I_1 = \int_0^1 x^{1+k+m+a_1+a_2} (1-x)^{b_1+b_2-l-n-1} \exp\{(\lambda_1 + \lambda_2)x\} dx,$$

$$I_2 = \int_0^1 x^{k+m+a_1+a_2} (1-x)^{b_1+b_2-l-n-1} \exp\{(\lambda_1 + \lambda_2)x\} dx$$

and

$$I_3 = \int_0^1 x^{k+m+a_1+a_2-1} (1-x)^{b_1+b_2-l-n-1} \exp\{(\lambda_1 + \lambda_2)x\} dx.$$

On applying (15), we can reduce these three integrals to:

$$I_1 = B(b_1+b_2-n-l, 2+m+k+a_1+a_2) \\ \times {}_1F_1(2+m+k+a_1+a_2; 2+k-l+m-n+a_1+b_1+a_2+b_2; \lambda_1+\lambda_2),$$

$$I_2 = B(b_1+b_2-n-l, 1+m+k+a_1+a_2) \\ \times {}_1F_1(1+m+k+a_1+a_2; 1+k-l+m-n+a_1+b_1+a_2+b_2; \lambda_1+\lambda_2)$$

and

$$I_3 = B(b_1+b_2-n-l, m+k+a_1+a_2) \\ \times {}_1F_1(m+k+a_1+a_2; k-l+m-n+a_1+b_1+a_2+b_2; \lambda_1+\lambda_2),$$

respectively. Substituting these into (38), we obtain an expression for  $R$  that is a finite sum of confluent hypergeometric functions. In the particular case  $b_1 = b_2 = 1$ , (38) reduces to the simple expression:

$$R = \exp\{-(\lambda_1 + \lambda_2)\} \left[ \frac{\lambda_1}{1+a_1+a_2} {}_1F_1(1+a_1+a_2; 2+a_1+a_2; \lambda_1+\lambda_2) \right. \\ \left. + \frac{a_1}{a_1+a_2} {}_1F_1(a_1+a_2; 1+a_1+a_2; \lambda_1+\lambda_2) \right].$$

**8. Log Beta distribution.** As the name indicates,  $X_i$  are said to have the log beta distribution if  $\log X_i$  have the standard beta distribution. Clearly the support of  $X_i$  must be a finite positive interval, say  $0 < e_i \leq X_i \leq f_i$ . Then  $Y_i = (\log X_i - \log e_i)/(\log f_i - \log e_i)$  will have the standard beta distribution for

some parameters, say  $a_i$  and  $b_i$ . Using this relationship, the reliability in (1) can be written as

$$\begin{aligned} R &= \Pr(X_2 < X_1) \\ &= \Pr\left(\frac{\log X_2 - \log e_2}{\log f_2 - \log e_2} < \frac{\log X_1 - \log e_2}{\log f_2 - \log e_2}\right) \\ &= \Pr\left(\frac{\log X_2 - \log e_2}{\log f_2 - \log e_2} < \frac{\log f_1 - \log e_1}{\log f_2 - \log e_2} \frac{\log X_1 - \log e_1}{\log f_1 - \log e_1} + \frac{\log e_1 - \log e_2}{\log f_2 - \log e_2}\right) \\ &= \Pr\left(Y_2 < \frac{\log f_1 - \log e_1}{\log f_2 - \log e_2} Y_1 + \frac{\log e_1 - \log e_2}{\log f_2 - \log e_2}\right). \end{aligned}$$

Hence, the reliability of the log beta distribution is the same as that of the generalized beta distribution given in (32) for the particular choices  $c_1 = (\log e_1 - \log e_2)/(\log f_2 - \log e_2)$ ,  $d_1 = (\log f_1 - \log e_2)/(\log f_2 - \log e_2)$ ,  $c_2 = 0$  and  $d_2 = 1$ . The log beta distribution has applications to the evolution of aerosol growth (Bunz et al., [1]; Chang et al., [2]).

**Acknowledgments.** The author is grateful to Professor Samuel Kotz (George Washington University, USA) and to Professor Marianna Pensky (University of Central Florida, USA) for introducing him to this area of research. The author is also grateful to a referee for the most useful comments and suggestions.

#### REFERENCES

- [1] C. BUNZ, W. SCHOCK, M. KOYRO, J. GENTRY, S. PLUNKETT, M. RUNYAN, C. PEARSON, C. WANG. Application of the log beta-distribution to aerosol size distributions. *Journal of Aerosol Science* **18**, (1987) 663–666.
- [2] Y. CHANG, R. J. HAN, C. L. PEARSON, M. R. RUNYAN, M. B. RANADE, J. W. GENTRY. Applications of the log beta-distribution to the evolution of aerosol growth. *Journal of Aerosol Science* **19**, (1988) 879–882.
- [3] I. S. GRADSHTEYN, I. M. RYZHIK. Tables of Integrals, Series, and Products. Academic Press, San Diego, 2000.
- [4] C. V. KIMBALL, D. J. SCHEIBNER. Error bars for sonic slowness measurements. *Geophysics* **63**, (1998) 345–353.



- [5] Y. W. LEE. Statistical Theory of Communications. John Wiley and Sons, New York, 1960.
- [6] M. MENICONI, D. M. BARRY. The power function distribution: A useful and simple distribution to assess electrical component reliability. *Microelectronics and Reliability* **36**, (1996) 1207–1212.
- [7] D. MIDDLETON. Introduction to Statistical Communications Theory. McGraw-Hill Book Company, New York, 1960.
- [8] W. L. NICHOLSON. A computing formula for the power of the analysis of variance test. *Annals of Mathematical Statistics* **25**, (1954) 607–610.
- [9] A. P. PRUDNIKOV, Y. A. BRYCHKOV, O. I. MARICHEV. Integrals and Series (volumes 1, 2 and 3). Gordon and Breach Science Publishers, Amsterdam, 1990.
- [10] B. ROUSSEAU, D. M. ENNIS. A Thurstonian model for the dual pair (4IAX) discrimination method. *Perception and Psychophysics* **63**, (2001) 1083–1090.
- [11] W. SHEN, P. D. ZHAO. Theoretical study of statistical fractal model with applications to mineral resource prediction. *Computer and Geosciences* **28**, (2002) 369–376.

*Department of Mathematics*  
*University of South Florida*  
*Tampa, Florida 33620, USA*  
*e-mail: snadaraj@chumal.cas.usf.edu*

*Received August 1, 2002*